

- Date: February 17, 2004
- Products: Dual II MAGNUM<sup>®</sup> Herbicide (EPA Reg No. 100-818)  
Dual II MAGNUM<sup>®</sup> SI Herbicide (EPA Reg No. 100-829)  
Dual IIG MAGNUM<sup>®</sup> Herbicide (EPA Reg No. 100-910)  
Dual MAGNUM<sup>®</sup> Herbicide (EPA Reg No. 100-816)  
Bicep II MAGNUM<sup>®</sup> Herbicide (EPA Reg No. 100-817)  
Bicep Lite II MAGNUM<sup>®</sup> Herbicide (EPA Reg No. 100-827)  
Liberty<sup>®</sup> Herbicide (EPA Reg No. 264-660)  
Callisto<sup>®</sup> (EPA Reg No. 100-1131)
- Use: Liberty at 20 oz/A may be applied post-emergence following a full labeled rate of Dual MAGNUM or Bicep MAGNUM brand herbicides applied at or before planting for improved control of Woolly Cupgrass in LibertyLink<sup>®</sup> corn. Callisto at 3 fl oz/Acre may be applied in tank mixture with Liberty for post-emergence broadleaf weed control.
- States: Illinois, Iowa, Minnesota, South Dakota, and Wisconsin

### **DIRECTIONS FOR USE**

It is a violation of federal law to use these products in a manner inconsistent with their labeling.

For improved control of woolly cupgrass, apply Liberty at 20 oz/A post-emergence following a full labeled rate of a Dual MAGNUM or Bicep MAGNUM brand herbicide applied at or before planting. Liberty must be applied to woolly cupgrass prior to tillering and when cupgrass is less than 3 inches in height. Apply Liberty in 20-30 gallons of water per acre. Callisto at 3 fl oz/A per acre may be tank mixed with Liberty for improved postemergence broadleaf weed control. Always add spray grade AMS at 3 lbs./A to the spray solution when applying Liberty alone (see the Liberty label for additional adjuvant instructions). AMS at 3 lbs./A plus a non-ionic surfactant at 0.25% v/v (1 qt/100 gal of water) is recommended when Callisto is applied in a tank mix combination with Liberty. No other adjuvants may be used with Liberty and Callisto tank mix combinations.

Notes: 1) Liberty may only be applied to corn specifically designated as having the LibertyLink trait. 2) Liberty may be applied to LibertyLink corn up to 24 inches tall or in the V-7 stage of growth. 3) Do not apply Liberty within 60 days of harvesting corn forage and within 70 days of harvesting corn grain and corn fodder.

**Refer to the Callisto and Liberty EPA approved container labels for complete directions for use and all applicable restrictions and precautions.**

**This labeling should be in the possession of the user at the time of pesticide application.**

**This recommendation is made as permitted by Section 2(ee) of FIFRA, as amended, and has not been submitted to or approved by EPA.**

Syngenta Crop Protection, Inc.  
Greensboro, NC 27419

Dual II MAGNUM<sup>®</sup>, Dual II MAGNUM<sup>®</sup> SI, Dual IIG MAGNUM<sup>®</sup>, Dual MAGNUM<sup>®</sup>, Bicep II MAGNUM<sup>®</sup>, Bicep Lite II MAGNUM<sup>®</sup> and Callisto<sup>®</sup> are trademarks of a Syngenta Crop Protection, Inc..

Liberty and LibertyLink are registered trademarks of Bayer Cropscience.

IL1131022AC0204

ILCallistoLibertyLinkCornSec2ee 2-17-04