

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Special 2(ee) Recommendation[†] (For Distribution and Use in the State of California)

**Aid in the Control of Certain Canker Diseases Following Pruning of Almonds, Apricots, Cherries,
Nectarines, Peaches, Plums and Prunes**

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Rally 40WSP according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Directions for Use

To protect pruning cuts from canker diseases caused by fungal pathogens such as *Eutypa lata*, **Leucostoma persoonii (Cytospora)**, **Calosphaeria pulchella**, **Lasiodiplodia theobromae**, and for suppression of *Botryosphaeria* spp. and **Diaporthe neotheicola (Phomopsis)**, apply Rally 40WSP fungicide at 6 oz per acre in 50 to 100 gallons of water per acre, using power operated ground application equipment or a backpack sprayer. If a backpack sprayer is used, chemical-resistant coveralls or a chemical-resistant coat must be worn. Lower application volumes per acre may be used only if the spray thoroughly wets all susceptible tissue.

For improved management of *Botryosphaeria* and *Diaporthe* and for fungicide resistance management, it is recommended to apply Rally 40WSP in a tank mix with Topsin M WSB at 1.5 lb/acre and/or in tank mix with other fungicides registered for canker disease management.

Protect pruning cuts as soon as possible, especially if irrigation, rainfall or fog is expected. Application within 24 hours is highly recommended. Applications after 24 hours are permitted, but disease control may be reduced. Assure thorough coverage of all cut wood surfaces. For these pathogens, multiple applications generally are needed. Make additional applications at approximately two week intervals, especially if irrigation, rainfall or high humidity conditions exist.

The addition of a labeled rate of a registered organosilicone spray adjuvant may increase penetration of cut wood surfaces. It is responsibility of the user to assure that the organosilicone spray adjuvant is safe to the crop under the existing conditions of use. Addition of a registered spray dye to the tank mix, and visual inspection of pruning cuts after application, will help to assure thorough coverage of all susceptible tissue.

Specific Use Restrictions:

Almonds: Do not apply within 90 days of harvest. Do not make more than 3 applications or apply more than a total of 1.5 lb of Rally 40WSP (0.6 lb ai) per acre per season.

Apricots, Plumbs, and Prunes: Applications may be made up to the day of harvest. Do not apply more than a total of 2.75 lb of Rally 40WSP (1.1 lb ai) per acre per season.

Cherries, Nectarines, Peaches: Applications may be made up to the day of harvest. Do not apply more than a total of 3.25 lb of Rally 40WSP (1.3 lb ai) per acre per season.

®Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

R182-066
Issued: 09/22/15
Initial printing.