

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 949A-S3 0615**

**Palisade® EC**

**This supplemental label expires on June 30, 2018 and must not be used or distributed after this date.**

<i>Active Ingredient:</i>	
Trinexapac-ethyl* .....	12.0%
<hr/>	
<i>Other Ingredients:</i>	88.0%
Total:	100.0%

\*CAS No. 95266-40-3

Palisade EC is an emulsifiable concentrate containing 1 pound of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**EPA Reg. 100-949**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Palisade EC as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

### Plant-back Interval (PBI):

Crop	Days to Plantback After the Last Application of Palisade EC
Wheat Barley Triticale Oats Rye Rice Grasses Grown for Seed Sugarcane	0 days
All Other Food or Feed Crops	30 days

Crop	fl oz/A (lb ai/A)	Application															
For yield protection and lodging prevention in rice																	
<b>Rice</b>  <b>Including wild rice</b>	3.5 – 5.7 (0.027 – 0.045)	<p>Apply Palisade EC one time as a broadcast spray to actively growing rice at the stages specified in the table below. Palisade EC may be applied via aerial or ground boom application (depending on when flooding takes place).</p> <p>Apply Palisade EC (one application only) within the following growth stages.</p> <table border="1"> <thead> <tr> <th>Growth Stage</th><th>BBCH code</th><th>Palisade EC Treatment</th></tr> </thead> <tbody> <tr> <td>Fully tillered</td><td>29</td><td>Yes (earliest)</td></tr> <tr> <td>Panicle Initiation (green ring)</td><td>30</td><td>Yes- Best</td></tr> <tr> <td>Panicle Formation (panicle 1-2mm)</td><td>32</td><td>Yes (latest)</td></tr> <tr> <td>Panicle Differentiation (jointing)/ ½- to ¾-inch internode elongation</td><td>34</td><td>NO – do not apply</td></tr> </tbody> </table> <p>Use the high rate on heavy, lush stands and varieties prone to lodging. Use the lower rate range when conditions are less favorable for lodging.</p> <p><b>NOTE:</b> Delayed heading has been noted when Palisade was applied later than panicle differentiation. As rice grows quickly, do not apply once internode is more than ½ - ¾ inch in length.</p>	Growth Stage	BBCH code	Palisade EC Treatment	Fully tillered	29	Yes (earliest)	Panicle Initiation (green ring)	30	Yes- Best	Panicle Formation (panicle 1-2mm)	32	Yes (latest)	Panicle Differentiation (jointing)/ ½- to ¾-inch internode elongation	34	NO – do not apply
Growth Stage	BBCH code	Palisade EC Treatment															
Fully tillered	29	Yes (earliest)															
Panicle Initiation (green ring)	30	Yes- Best															
Panicle Formation (panicle 1-2mm)	32	Yes (latest)															
Panicle Differentiation (jointing)/ ½- to ¾-inch internode elongation	34	NO – do not apply															

### Specific Use Restrictions:

- 1) Do not apply more than 5.7 fl oz (0.045 lb ai/A) Palisade EC/A/year.
- 2) Do not apply to ratoon crop.
- 3) Do not release flood water for 5 days after application.
- 4) Do not apply after 50 days pre-harvest (50-day PHI).

Palisade® and the Syngenta logo are trademarks of a Syngenta Group Company

©2015 Syngenta