

Bayer CropScience LP P.O. Box 12014 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937) http://www.bayercropscienceus.com Baythroid<sup>®</sup> XL

EPA Reg. No. 264-840

Use Directions for:

Brassica (COLE) Leafy Vegetables Crop Group 5 and Leafy Vegetables Crop Group 4

# PRODUCT BULLETIN

## RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

## FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

### **BRASSICA (COLE) LEAFY VEGETABLES**

### Includes all members of Crop Group 5 such as, but not limited to:

Broccoli, Broccoli raab (rapini); Chinese (gai lon) broccoli; Brussels sprouts; Cabbage; Chinese (bok choy) cabbage; Chinese (napa) cabbage; Chinese mustard (gai choy) cabbage; Cauliflower; Cavalo broccoli; Collards; Kale; Kohlrabi; Mizuna; Mustard greens; Mustard spinach; Rape greens; and Turnip greens.

#### **LEAFY VEGETABLES**

Includes all members of Crop Group 4 such as, but not limited to: Amaranth (Chinese spinach); Arugula (rouquette); Cardoon; Celery; Chinese celery; Celtuce; Chervil; Chrysanthemum (edible-leaved and garland); Corn salad; Cress (garden and upland); Dandelion; Dock (sorrel); Endive (escarole); Florence fennel; Lettuce (head and leaf); New Zealand spinach; Orach; Parsley; Purslane (garden and winter); Radicchio, (red chicory); Rhubarb; Spinach; Swiss chard; Vine spinach.

PESTS CONTROLLED	Rate fluid ounces/Acre	Rate Ib Al/Acre	Method
Cabbage Maggot Garden Symphylan Seed Corn Maggot	3.2 fl.oz. per acre	0.025	For reduction of damage caused by the listed pests, apply specified dosage per acre as a soil-incorporated treatment before or at planting or transplanting.

#### Notes and Restrictions:

Maximum number of pre-plant or at-plant/transplant applications: 1.

Maximum BAYTHROID XL allowed per crop season: 12.8 fl.oz. (0.1 lb Al/Acre).

See Chemigation Statement in Application Guidelines on the specimen label

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions. For additional information call toll-free 1-866-99BAYER (1-866-992-2937)