



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

SILVERADO™ Wild Oat Herbicide

EPA Reg. No. 264-801

OLYMPUS™ Herbicide

EPA Reg. No. 264-809

Silverado Wild Oat herbicide plus OLYMPUS herbicide in tank mixture for use on wheat to control certain emerged annual grasses and broadleaf weeds.

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF MINNESOTA, MONTANA, NORTH DAKOTA, SOUTH DAKOTA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Product	Rate	Crop	Weeds
OLYMPUS™ Herbicide + Silverado™ Wild Oat Herbicide + MSO or Basic Blend type adjuvant	0.2 ounce/acre + 2.75 ounces/acre + 1.5 pt MSO/acre or 1% v/v Basic Blend adjuvant	Apply tankmixture to wheat from emergence up to jointing stage of growth	For control of wild oats, yellow foxtail*, green foxtail*, Persian darnel*, barnyardgrass, true cheat*, Japanese brome*, foxtail barley*, mustards and volunteer canola

*These weeds will be controlled when 2.75 oz / Acre Silverado + 0.2 oz / Acre Olympus are applied in tankmixture with a recommended adjuvant system at the pre-tiller stage of weed growth.

Partially Controlled Weeds

Grass weeds that will be partially controlled: Downy brome, smooth brome, little barley and quackgrass.

Broadleaf weeds that will be partially controlled: Catchweed bedstraw, common chickweed, cornflower / bachelor's button, field pennycress, henbit, ivyleaf speedwell and mouseear chickweed.

Rotational Crop Interval for SILVERADO Wild Oat herbicide Tank Mixed with Olympus

Up to 0.2 oz/A of Olympus 70 WG /acre may be tank mixed with SILVERADO Wild Oat herbicide.

CROP	ROTATION INTERVAL Months
Wheat	0
Millet	4
Alfalfa	10
Barley	10
Canola	10
Dry beans	10
Flax	10
Lentils	10
Peas	10
Safflower	10
Soybeans	10
Sugar beets	10
Sunflowers	10
Corn	12

24 inches of precipitation and a 12 month rotation interval are required for potatoes, buckwheat, and onions.

DIRECTIONS FOR USE

- A recommended adjuvant must be tankmixed according to the guidelines on the federal Silverado Wild Oat herbicide label.
- Applications should be made to actively growing annual grasses and broadleaf weeds. Weed control may be reduced when weeds are under stress due to severe weather conditions, drought, very cold temperatures, etc. Weed control may be reduced if the herbicide application is made under dry, dusty conditions – especially in the wheel track areas.
- Do not apply when wind causes drift to off-site vegetation as injury may occur.

For use directions, precautions and restrictions not addressed in this bulletin follow the most restrictive directions on the respective labels of both partners in the tank mixture.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).

OLYMPUS and Silverado Wild Oat herbicide are trademarks of Bayer.