



**Bayer CropScience**

Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)

## Poncho® 600

Add Sugarcane Aphid (*Melanaphis sacchari*)  
for Commercial Sorghum Seed Treatment.

EPA Reg. No. 264-789

## FIFRA 2(ee) Recommendation

### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF ALABAMA, ARKANSAS, FLORIDA, GEORGIA, KANSAS, LOUISIANA, MISSOURI, MISSISSIPPI, NEBRASKA, NEW MEXICO, OKLAHOMA, TENNESSEE AND TEXAS**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its label.

#### **COMMERCIAL SORGHUM SEED TREATMENT**

Pest	Rate of Poncho 600	Remarks
Sugarcane Aphid	Apply 5.1 to 6.4 fl oz per 100 lbs seed	Apply prior to planting as a slurry treatment to provide early season protection of seedlings. Ensure thorough coverage.

#### **SEED BAG TAG**

The Federal Seed Act requires that the container of seed treated with Poncho 600 must be labeled with the following statements:

- This seed has been treated with Poncho 600, which contains clothianidin.
- Do not use treated seed for food, feed or oil processing.

In addition, the US Environmental Protection Agency requires the following statements on the container of seed treated with Poncho® 600:

- Store away from feeds and foodstuffs.
- Rapeseed greens and seed grown or harvested from Poncho 600-treated seed must not be used for feed or human consumption. Rapeseed grown and harvested from Poncho 600-treated seed is only for industrial uses and cannot be used for edible oil or any other human/feed consumption.
- Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves when handling treated seed.
- Dispose of all excess treated seed. Left over treated seed may be double sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- This compound is toxic to birds and mammals. Treated seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect treated seeds spilled during loading.
- For Sweet corn, Canola, Rapeseed, and Sorghum: Immediate Plant-back: Cereal grains, Field corn, popcorn, sweet corn, rapeseed, soybeans, canola, sorghum, root and tuber vegetables, 30-Day Plant-back: Grasses, non-grass animal feeds, dried beans, 4-Month Plant-back: Any crop without an earlier plant-back interval

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.  
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937).

*As with any crop-protection product, always read and follow label directions.  
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*