



Bayer CropScience LP
P.O. Box 12014,
2 T.W. Alexander Drive
Research Triangle Park, NC 27709
(919) 549-2000

PROLINE® 480 SC Fungicide

EPA Reg. No. 264-825

For Suppression of Sclerotinia Stem Rot
(*Sclerotinia sclerotiorum*) in Soybean

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NEW YORK

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

SOYBEAN	
Disease Suppressed	Rate fluid ounces/Acre
Sclerotinia Stem Rot also known as White Mold (<i>Sclerotinia sclerotiorum</i>)	3.0
Notes: PROLINE® 480 SC FUNGICIDE may be applied by either ground, aerial or chemigation application equipment. For aerial application, apply in a minimum spray volume of 2 gpa. Apply PROLINE® 480 SC FUNGICIDE as a broadcast, preventative foliar spray or at first visible symptoms of the disease. Repeat applications on a 10- to 21-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rate and shorter spray intervals are recommended when disease pressure is severe.	
Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 9 fl oz. of PROLINE® 480 SC FUNGICIDE per acre per use season.	

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).