## THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA



EPA Reg. No. 71512-3-279

**Group 21 Fungicide** 

For Agricultural or Commercial Use Only

## 2(ee) Recommendation FOR DISTRIBUTION AND USE ONLY IN

California, Colorado, Delaware, Idaho, Kansas, Maine, Maryland, Michigan, Montana, Nebraska, Nevada, New Jersey, North Carolina, Oregon, Pennsylvania, Utah, Virginia, Washington, and West Virginia

Virginia, Washington, and West Virginia
THIS PRODUCT BULLETIN CONTAINS NEW OR
ADDITIONAL DIRECTIONS FOR USE WHICH ARE
RECOMMENDED BY FMC CORPORATION UNDER
FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

Сгор	Pest	Rate of Application
Potatoes	Pythium ultimum suppression	In-furrow at plant: 0.42fluid oz per 1000 linear feet (Equivalent to 6.1 fl. oz. per Acre on 36 inch row spacing) Lay by/Hilling: 2.75 fl. oz. per Acre

## **DIRECTIONS FOR USE:**

Thorough coverage is essential to achieve control. Apply with sufficient spray volume to obtain full coverage of the furrow, at planting prior to furrow coverage.

**Potatoes (7 Day PHI):** For Pythium ultimum suppression, apply at planting, 0.42 fluid ounces per 1000 linear feet (Equivalent to 6.1 fluid oz/acre on 36 inch row spacing) of row in-furrow at planting using a minimum of 5 gallons of water per acre. Apply RANMAN FUNGICIDE using a 6 to 8 inch band directly over the seed pieces prior to furrow closure. A side dressing of RANMAN FUNGICIDE applied at hilling may be necessary for additional control. Where mefenoxam-resistant strains of Pythium ultimum are not present, a full rate of RANMAN FUNGICIDE can be tank-mixed with mefenoxam containing fungicide for additional control.

For additional suppression of Pythium ultimum in combination with an at-planting, in-furrow RANMAN FUNGICIDE application, apply RANMAN FUNGICIDE as a broadcast spray at 2.75 fl. oz per Acre in a minimum of 20 gallons of finished spray solution per acre at hilling. Additional applications may be needed depending on susceptibility of potato varieties to Pythium spp., environmental conditions conducive to favor severe disease development, or fields in long growing season areas, etc.

## **RESTRICTIONS:**

Do not apply more than 27.5 fluid ounces product per acre per year.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED RANMAN FUNGICIDE LABEL (EPA REG. NO. 71512-3-279) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



