



# SUPPLEMENTAL LABEL

## Mycoshield®

FOR THE CONTROL OF FIRE BLIGHT OF PEAR AND APPLE AND  
BACTERIAL SPOT OF PEACH AND NECTARINE

EPA Reg. No. 55146-97  
06-21-2012

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR THIS PRODUCT BEFORE  
PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.  
"Label" as used in this supplemental labeling refers to the label booklet for this product and this supplemental label.

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING.  
This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS IN THE PRODUCT LABEL BOOKLET MUST BE FOLLOWED, INCLUDING STATEMENTS PERTAINING TO THE WORKER PROTECTION STANDARDS, ON THE EPA REGISTERED LABEL.

### APPLE and PEAR

DISEASE CONTROLLED	CONCENTRATION / USE RATE	APPLICATION INSTRUCTIONS
<b>Fire Blight</b> ( <i>Erwinia amylovora</i> )	<b>Ground Application:</b> 200 ppm Equivalent to: 1.0 lb Mycoshield / 100 gal water	<b>Ground Application:</b> Apply using airblast sprayer. Ensure good coverage of plant parts. Apply beginning with 10% bloom. <u>For Apples</u> , repeat applications at 3 to 6 day intervals or apply when blight favorable weather is expected during apple bloom. <u>For Pears</u> , repeat applications at 4 to 6 day intervals.
	<b>Aerial Application:</b> 1.0 lb Mycoshield / minimum 10 gal water	<b>Aerial Application:</b> Application by air is not a direct replacement for applications by ground. Use aerial application only in those orchards in which optimal coverage can be achieved by air. Even with optimal coverage, the extent of control may be reduced. Apply by ground within 3 to 6 days subsequent to an aerial application. Begin spray application at 10% bloom. <u>For apples</u> , repeat applications at 3 to 6 day intervals or apply when blight favorable weather is expected during apple bloom. <u>For pears</u> , repeat applications at 4 to 6 day intervals. Spray coverage of blossoms is essential.

#### Restrictions:

##### For Apples:

DO NOT apply more than 1.5 lb of product per acre per application.  
DO NOT exceed a total of 6 applications per year using either ground or aerial methods of application.  
DO NOT apply within 60 days of harvest.

##### For Pears:

Do not apply more than 1.0 lb of product per acre per application.  
DO NOT exceed a total of 10 applications per year using either ground or aerial methods of application.  
DO NOT apply within 60 days of harvest.

**NOTE:** Use of this product may cause phytotoxicity to the fruit and/or foliage of sensitive pear varieties, especially Asian varieties.

### NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Manufactured for NUFARM AMERICAS INC. AGT DIVISION  
150 Harvester Drive, Burr Ridge, IL 60527

Mycoshield is a trademark of Nufarm Americas Inc.