

## SUPPLEMENTAL LABEL

# Confine<sup>®</sup> Extra

## Fungicide

EPA Reg. No. 1381-244

### KEEP OUT OF REACH OF CHILDREN

## CAUTION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read this label and the product container label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for Confine Extra attached to the container.

### CEREAL GRAINS, FORAGE AND FIBER CROP GROUPING

Cereal Grains, Forage and Fiber Crop Grouping: Alfalfa, Barley, Buckwheat, Canola, Clover (all types), Coastal Bermudagrass, Corn (all types), Cotton, Hay, Kudzu, Lespedeza, Lupin, Millet, Oats, Popcorn, Rice, Rye, Safflower, Sainfoin, Sorghum, Sunflower, Teosinte, Trefoil, Triticale, Vetch, Wheat and Wild rice.

Make applications prior to disease development in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Use rates and frequency are designed to prevent plant injury. If more frequent applications are necessary for disease control, consult a crop expert and follow their recommendations. Do not exceed the highest rate per application. Do not apply at less than 3 day intervals. Do not apply to plants that are dormant or heat or moisture stressed. To avoid undesirable copper phytotoxicity, do not apply CONFINE EXTRA FUNGICIDE foliarly to plants treated with copper based compounds at less than 20 day intervals unless instructed to do so by your consultant. Allow foliage to completely dry after application. Do not apply when conditions favor wet tissue for prolonged periods (>4 hours).

Disease	Application Method	Rate	Application Program
Phytophthora Pythium Fusarium Rhizoctonia	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Disease Prevention Program*: Apply lower rate at 2-4 week intervals after plants become established. Do not apply CONFINE EXTRA FUNGICIDE at intervals less than 3 days. Disease Control Program: Apply higher rate at 2-3 week intervals until control is reached. Consult with Farm Advisor or crop expert to determine disease severity. Do not apply CONFINE EXTRA FUNGICIDE at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	
	Irrigation	Overhead: 2-4 quarts in at least 100 gallons of water.  Drip: 2-3 quarts in at least 100 gallons of water.	Apply with normal irrigation schedule. Follow Disease Prevention Program or Disease Control Program as stated above. Do not apply CONFINE EXTRA FUNGICIDE at intervals less than 3 days.
Downy Mildew	Foliar	1 to 3 quarts in a minimum of 20 gallons of water/acre.	Apply at 2-3 week intervals. Follow Disease Prevention Program or Disease Control Program as stated above. Do not apply CONFINE EXTRA FUNGICIDE at intervals less than 3 days.
	or  Aerial Application	1 to 3 quarts in a minimum of 10 gallons of water/acre.	

\*Check with your local Extension Agent if you are unsure these conditions exist.

CONFINE<sup>®</sup> is a registered trademark of Winfield Solutions, LLC.

DISTRIBUTED BY:  
WINFIELD SOLUTIONS, LLC  
P.O. Box 64589  
St. Paul, MN 55164-0589