

Carnivore™ Herbicide

FIFRA Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY
IN THE STATE OF NEBRASKA**

FOR POSTEMERGENCE CONTROL OF KOCHIA ON FALLOWLAND

ACTIVE INGREDIENTS:

MCPA-EHE: 2-methyl-4-chlorophenoxyacetic acid, 2-ethylhexyl ester* 27.30%

Octanoic acid ester of bromoxynil (3,5-dibromo-4-hydroxybenzonitrile)** 25.48%

Fluroxypyr 1-methylheptyl ester ((4-amino-3,5-dichloro-6-fluoro-2-pyridinyl)oxy) acetic acid, 1-methylheptyl ester*** 10.08%

OTHER INGREDIENTS: 37.14%

TOTAL: 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This label and the Federal label for this product must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Carnivore Herbicide and this Section 24(c) label before applying. Do not use in a manner inconsistent with applicable use directions, restrictions, precautions and limitations provided on the label affixed to the container for Carnivore Herbicide and this Section 24(C) label.

USE ON FALLOWLAND

| Weed | Rate of Application | Comments |
|--------|-----------------------|---|
| Kochia | 1 to 1-1/2 pints/Acre | <p>Spray Volume: Use sufficient spray volume to provide thorough coverage and a uniform spray pattern. For best results (and to minimize spray drift), apply in a spray volume of 8 gallons of water or more per acre by ground and 3 or more gallons of water per acre by air. Spray volume should be increased as weed density and vegetative canopy increase in order to obtain equivalent weed control; however, do not exceed 40 gallons per acre total spray volume.</p> <p>Notes re Kochia: Research indicates many biotypes of kochia may occur within a single field and while kochia biotypes can vary in their susceptibility to this product, in general all biotypes will be suppressed or controlled at the labeled rate. A shift to more tolerant biotypes within a field may occur if this product is applied at rates lower than recommended.</p> <p>Kochia seedlings less than 4 inches tall (including ALS resistant biotypes) will be controlled using the 1 pint per acre rate. However, when conditions for control are less favorable, such as under drought or cool temperature, a rate of 1-1/2 pints per acre will provide more consistent control of kochia seedlings 1 to 4 inches tall. For more consistent control of small kochia, apply when the plants are at least 1 inch tall. A rate of 1-1/2 pints per acre should be used for optimal control of dicamba tolerant kochia populations.</p> <p>Restrictions: Do not allow livestock to graze in treated areas or feed treated grass to livestock.</p> |

EPA Reg. No. 1381-249

EPA SLN No. NE-140002

Distributed By
Winfield Solutions, LLC
P.O. Box 64589, St. Paul, MN 55164-0589

WINFIELD™

1/0409/4