

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 828A-S6 0915**

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>9</b> | FUNGICIDE |
|-------|----------|-----------|

**Vangard® WG**

### Fungicide

**This supplemental label expires on 09/30/2018 and must not be used or distributed after this date.**

|   |        |
|---|--------|
| Active Ingredient:  |        |
| Cyprodinil: 4-cyclopropyl-6-methyl-N-phenyl-pyrimidinamine* ..... | 75.0%  |
| Other Ingredients:  | 25.0%  |
| Total:  | 100.0% |

\*CAS No. 121552-61-2

Vangard WG is a water-dispersible granule.

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

**EPA Reg. No. 100-828**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Vangard WG as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

### Rotational Crop Restrictions

Do not plant any crop which is not registered for use with cyprodinil for a period of 30 days, unless a shorter interval is specified on the following list.

| Rotation Crop   | Planting Time From Last Vanguard WG Application |
|---|---|
| Artichoke, Globe<br>Beans (dried and succulent except cowpeas)<br>Berries (bushberries 13-07B, caneberries 13-07A)<br>Brassica (Cole) Leafy Vegetables<br>Cucurbits<br>Herbs (fresh and dried)<br>Leafy Vegetables<br>Leaves of Root and Tuber Vegetables<br>Onions (dry bulb, including garlic, and green)<br>Peppers<br>Root Vegetables except Sugar beet<br>Strawberries<br>Tomatoes and tomatillos<br>Watercress<br>Crops Not Intended for Food or Feed | 0 days  |
| All Other Crops Intended for Food or Feed   | 30 days   |

**Restriction:** In annual crops, where multiple crops can be grown per year (double/triple cropping), do not apply more than 1.3 lb ai per acre per year to an individual plot of land.

| Crop  | Disease   | Product Rate oz/Acre                      | Remarks   |
|---|---|---|---|
| <b>Stone Fruits</b><br><b>Crop Group 12-12*</b><br><br>Apricots<br>Tart cherries<br>Nectarines<br>Peaches<br>Plums<br>Prunes<br><br>And cultivars and/or hybrids of these | Brown rot blossom blight<br>( <i>Monilinia</i> spp.)                      | 5   | Begin applications at bloom stage (apricots at red bud, cherries at popcorn, peaches and nectarines at pink bud, plums and prunes at green tip). Make a second application at full bloom. Use Vanguard WG alone or in tank mixture with the label rate of another fungicide registered for stone fruit. |
|   | <b>California only</b><br><br>Fruit brown rot<br>( <i>Monilinia</i> spp.) | <b>Vanguard WG alone</b><br><br>10        | Apply a maximum of 2 applications of Vanguard WG during the preharvest period up through 2 days prior to harvest as needed.   |
|   |   | <b>Vanguard WG tank mixtures</b><br><br>5 | Apply Vanguard WG in tank mixture with the label rate of another fungicide registered on stone fruit for control of fruit brown rot.  |

**\*Complete List of Stone Fruit:** Apricot; Apricot, Japanese; Cherry, tart; Jujube, Chinese; Nectarine; Peach; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plum, prune; Plumcot; Prune; Sloe; as well as other cultivars and hybrids of these

| Crop | Disease | Product Rate<br>oz/Acre | Remarks |
|------|---------|-------------------------|---------|
|------|---------|-------------------------|---------|

**Stone Fruits**  
**Crop Group 12-12\***  
**(continued)**

**Application Instructions:** Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

**Specific Use Restrictions**

- 1) Do not apply Vangard WG to sweet cherries.
- 2) Make no more than two applications by air.
- 3) Do not apply more than 30 oz/A of Vangard WG (1.4 lb ai/A of cyprodinil) per year.
- 4) Do not apply more than a maximum total of 4 applications (air plus ground) per year.
- 5) Do not apply within 2 days of harvest (2-day PHI).

| Crop                   | Disease   | Product Rate<br>oz/Acre | Remarks   |
|------------------------|---|-------------------------|---|
| <b>Tropical Fruits</b> | Botrytis fruit rot<br>( <i>Botrytis</i> spp.)     | 5.5 - 7                 | Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development.      |
| Acerola                | Alternaria fruit rot<br>( <i>Alternaria</i> spp.) |                         | Resistance Management: After 2 applications of Vangard WG, alternate with another fungicide with a different mode of action for 2 applications. |
| Avocado                |   |                         |   |
| Black sapote           |   |                         |   |
| Canistel               |   |                         |   |
| Dragon Fruit           | Anthracnose<br>( <i>Colletotrichum</i> spp.)      |                         |   |
| Feijoa                 |   |                         |   |
| Guava                  |   |                         |   |
| Jaboticaba             |   |                         |   |
| Lychee                 |   |                         |   |
| Longan                 |   |                         |   |
| Mamey sapote           |   |                         |   |
| Mango                  |   |                         |   |
| Papaya                 |   |                         |   |
| Passionfruit           |   |                         |   |
| Pulasan                |   |                         |   |
| Rambutan               |   |                         |   |
| Sapodilla              |   |                         |   |
| Spanish lime           |   |                         |   |
| Star apple             |   |                         |   |
| Starfruit              |   |                         |   |
| Wax jambu              |   |                         |   |

**Application Instructions:** Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.

**Specific Use Restrictions**

- 1) Make no more than two applications by air.
- 2) Do not apply more than 28 oz/A of Vangard WG (1.3 lb ai/A of cyprodinil) per year.
- 3) Do not apply more than a maximum total of 4 applications (air plus ground) per year.
- 4) May be applied on the day of harvest (0-day PHI).

Vangard® and the Syngenta logo are trademarks of a Syngenta Group Company

©2015 Syngenta