



# Supplemental Label

## METEOR™ Fungicide

EPA Reg. No. 70506-243

**This supplemental label expires on July 1, 2018, and must not be used or distributed after this date.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The supplemental labeling and the entire METEOR Fungicide container label, EPA Registration Number 70506-243, must be in the possession of the user at the time of the application. Read the label affixed to the container for METEOR Fungicide before applying. Carefully follow all precautionary statements and applicable use directions. Use METEOR Fungicide according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container for METEOR Fungicide.

### BERRIES

**Caneberry: Blackberry, loganberry, red and black raspberry, cultivars and/or hybrids of these.**

**Bushberry: Currant, elderberry, gooseberry, huckleberry**

**Blueberry: highbush and lowbush, uses allowed West of the Rocky Mountains only.**

DISEASE	RATE/ACRE	INSTRUCTIONS
<b>Botrytis Fruit Rot</b> ( <i>Botrytis cinerea</i> )	1.0 – 2.0 pts/A	Apply in a minimum of 100 gallons of water per acre. Apply at early bloom (5 to 10% bloom) and again at full bloom. Two additional applications may be made at 14 day intervals or as required. Use METEOR as an integral part of a complete disease control program. Apply as a foliar spray in sufficient water to ensure thorough coverage of both blossoms and fruit. Under severe disease conditions, use the high rate. For blueberry use West of the Rocky Mountains: METEOR may be sensitive to certain blueberry varieties. Test variety sensitivity prior to use.

### RESTRICTIONS

**METEOR is only registered for use on blueberries West of the Rocky Mountains. Do not use METEOR fungicide on blueberries East of the Rocky Mountains.**

Do not make more than 4 applications per season on bushberries (currant, elderberry, gooseberry, and huckleberry).

Do not make more than 4 applications per season on caneberries or blueberries, highbush and lowbush (West of the Rocky Mountains only).

The final application can be made up to and including the day of harvest (PreHarvest Interval (PHI)= 0 days)

**Rev. 07/01/2015**

United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
1-800-438-6071