



Supplemental Label

Approval pending in some states

MANZATE® FLOWABLE FUNGICIDE

EPA Reg. No. 70506-236

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. MANZATE® FLOWABLE Fungicide must be used only in accordance with the use directions on this label.

USE RATES AND APPLICATION TIMINGS

(See table on pages two and three attached)

IMPORTANT

BEFORE USING MANZATE® FLOWABLE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Read the Limitation of Warranty and Liability on the Section 3 Federal Product labeling before buying or using MANZATE® FLOWABLE. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Crops Included	Diseases Controlled	Rate per application Manzate® Flowable Qts/Acre	Remarks	Restrictions
Sugar apple Cherimoya Atemoya Custard apple Sweetsop	Anthraco nose	1.6-1.8	Begin applications at flowering and continue at a 7-day retreatment interval. Applications made with aerial equipment must be made in a minimum spray volume of 10 gal/acre.	Do not apply more than 26.25 qts. product (26.25 lbs. active ingredient) per acre per year. Do not apply more than 14 applications per year. Applications may be



Supplemental Label

Crops Included	Diseases Controlled	Rate per application Manzate® Flowable Qts/Acre	Remarks	Restrictions
				made up to the day of harvest.
Ginseng	Alternaria blight	1.5	<p>Start applications when disease first threatens and repeat every 7-10 days as needed.</p> <p>In Wisconsin, apply with ground equipment and a minimum of 80 gallons of water per acre.</p>	<p>Do not apply more than 18 qts. product (18 lbs. active ingredient) per acre per year.)</p> <p>Do not apply more than 12 applications per year.</p> <p>Do not apply within 30 days of harvest.</p>
Cucurbit crop group Chayote Chinese wax gourd Citron melon Cucumber Gherkin Gourd, edible Momordica spp. Muskmelon Pumpkin Squash, summer Squash, winter Watermelon	Anthracnose Cercospora leaf spot Downy mildew Gummy stem blight Scab Alternaria leaf spot	1.6-2.4	<p>Start applications when the plants are in the two-leaf stage and repeat at 7- to 10- day intervals.</p> <p>Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces.</p> <p>For aerial applications, the minimum spray volume is 2 gallons per acre.</p> <p>Some cantaloupe varieties (i.e. Harvest Queen, Gold Star, Super Star, Sweet and Early, and</p>	<p>Do not apply more than 19.2 qts. product (19.2 lbs. active ingredient) per acre per year.</p> <p>Do not apply more than 8 applications per year.</p> <p>Do not apply within 5 days of harvest.</p>



Supplemental Label

Crops Included	Diseases Controlled	Rate per application Manzate® Flowable Qts/Acre	Remarks	Restrictions
			Saticoy) are sensitive to Manzate® Flowable fungicide. Consult State Cooperative Extension Service Specialist prior to use.	
Mango Star apple (caimito) Canistel Mamey sapote Sapodilla White sapote	Anthracnose Phytophthora fruit rot Black spot (Cercospora)	1.6-2.0	Start applications at flowering and continue at 14- to 21-day intervals. Direct spray to crown and blossom area. Use 20 to 100 gallons water per acre.	Do not apply more than 28 qts. product (28 lbs. active ingredient) per acre per year. Do not apply more than 14 applications per year. Applications may be made up to the day of harvest.