



Supplemental Label

DEVIRINOL[®] 50-DF

Selective Herbicide - Dry Flowable
(EPA Reg. No. 70506-36)

SUPPLEMENTAL LABEL FOR USE ON CRANBERRIES

ACTIVE INGREDIENT:

Napropamide 50%

OTHER INGREDIENTS: 50%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This supplemental label must be in the possession of the user at the time of application. The use of Devrinol 50-DF according to this label is subject to the use precautions and limitations found on the EPA-registered label affixed to the Devrinol 50-DF container.

Read the entire Precautionary Statements, Directions for Use and Conditions of Sale and Limitation of Warranty and Liability found on the container label before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Always read and follow the label directions and any precautions or restrictions when applying any pesticide alone or in combination with tank mix partners.

CAUTION: Do not apply in the spring to cranberry beds that do not have sprinkler systems for frost control.

Northeast (Connecticut, Delaware, Massachusetts, Maryland, Maine, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont, West Virginia)

Established beds (over one year old)

For control of nutsedge (*Cyperus dentatus*) and rice cutgrass (*Leersia oryzoides*) and purplestem beggarticks (*Bidens connata*), apply DEVIRINOL 50-DF at the rates indicated below to a weed free soil surface before spring growth begins or in the fall after harvest. Do not apply when beds are under winter flood.

In Peat Beds: Apply 12-18 pounds DEVIRINOL 50-DF per acre through ground application equipment or through the overhead sprinkler system. See use directions for application through sprinkler irrigation systems. Do not apply more than 18 pounds DEVIRINOL 50-DF per acre per year.

In Beds with Sandy Soils: Apply 8-12 pounds DEVIRINOL 50-DF per acre through ground application equipment or through the sprinkler irrigation system. Do not apply more than 12 pounds DEVIRINOL 50-DF per acre per year.

New Plantings

Apply 6 pounds DEVIRINOL 50-DF per acre to a weed free soil surface after planting using ground spray equipment or through the sprinkler irrigation system. Do not apply more than 6 pounds DEVIRINOL 50-DF per acre per year.

North Central (Michigan, Minnesota, North Dakota, South Dakota, Wisconsin)

Established beds (over one year old)

For control of purplestem beggarticks (*Bidens connata*) and rice cutgrass (*Leersia oryzoides*), apply DEVRINOL 50-DF at the rates indicated below to a weed free soil surface before spring growth begins or in the fall after harvest. Avoid applying when beds may be reflowed for spring frost protection or when soils are water soaked. Do not apply when beds are under winter flood.

In Peat Beds: Apply 12-18 pounds DEVRINOL 50-DF per acre through ground application equipment or through the overhead sprinkler system. See use directions for application through sprinkler irrigation systems. Do not apply more than 18 pounds DEVRINOL 50-DF per acre per year.

In Beds with Sandy Soils: Apply 8-12 pounds DEVRINOL 50-DF per acre through ground application or through the sprinkler irrigation system. Do not apply more than 12 pounds DEVRINOL 50-DF per acre per year.

New Plantings

Apply 6 pounds DEVRINOL 50-DF per acre to a weed free soil surface after planting using ground spray equipment or through the sprinkler irrigation system. Do not apply more than 6 pounds DEVRINOL 50-DF per acre per year.

West (Idaho, Montana, Oregon, Washington)

Established beds (over one year old)

For control of clover (*Trifolium fimbriatum*), aster (*Aster douglasii*), silverweed cinquefoil* (silverleaf) (*Potentilla anserina*), red sorrel (sheep sorrel) (*Rumex acetosella*) and birdsfoot trefoil (*Lotus corniculatus*), apply DEVRINOL 50-DF at the rates indicated below to a weed free soil surface before spring growth begins in mid-February or before the end of March. Do not apply after buds start to swell.

In Beds with Heavy Muck Soil: Apply 18 pounds of DEVRINOL 50-DF per acre through ground application equipment or through the overhead sprinkler system. See use directions for application through sprinkler irrigation systems. Do not apply more than 18 pounds DEVRINOL 50-DF per acre per year.

In Peat Beds: Apply 12 to 18 pounds DEVRINOL 50-DF per acre through ground application equipment or through the overhead sprinkler system. Do not apply more than 18 pounds DEVRINOL 50-DF per acre per year.

In Beds with Sandy Soils: Apply 8-18 pounds DEVRINOL 50-DF per acre through ground application equipment or through the sprinkler irrigation system. Do not apply more than 18 pounds DEVRINOL 50-DF per acre per year.

New Plantings

Apply 6 pounds DEVRINOL 50-DF per acre to a weed free soil surface after planting using ground spray equipment or through the sprinkler irrigation system. Do not apply more than 6 pounds DEVRINOL 50-DF per acre per year.

*Suppression only

Rev. 11/07

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
www.upi-usa.com • 1-800-438-6071