



Supplemental Label

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

BIFENTURE® EC AGRICULTURAL INSECTICIDE

EPA Reg. No. 70506-57

NOTE: Use on grasses is limited to the States of Idaho, Oregon, and Washington.

This supplemental label expires on December 31, 2018, and must not be used or distributed after this date.

For use on Grass Forage, Fodder, and Hay Group and Grass Grown for Seed, Pasture and Rangeland (including: bahiagrass, barnyardgrass, bentgrass, Bermudagrass, Kentucky bluegrass, big bluestem, smooth brome grass, buffalograss, reed canarygrass, centipedegrass, crabgrass, cupgrass, dallisgrass, sand dropseed, Kentucky fescue, meadow foxtail, eastern gramagrass, side-oats grama, guinea grass, Indian grass, Johnsongrass, lovegrass, napiergrass, oatgrass, orchardgrass, pangolagrass, paspalum, redtop, Italian ryegrass, St. Augustine grass, sprangletop, squirreltailgrass, stargrass, switchgrass, timothy, crested wheatgrass, wildrye grass and zoysis grass. Also included are sudangrass and sorghum forages and their hybrids).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The supplemental labeling and the entire BIFENTURE EC container label, EPA Registration Number 70506-57, must be in the possession of the user at the time of the application. Read the label affixed to the container for BIFENTURE EC before applying. Carefully follow all precautionary statements and applicable use directions. Use BIFENTURE EC according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container for BIFENTURE EC.

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071



Supplemental Label

PESTS CONTROLLED	RATE BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS
Alfalfa Caterpillar Alfalfa Looper Alfalfa Weevil Blue Alfalfa Aphid ¹ Cutworms Egyptian Alfalfa Weevil (larvae & adult) Flea Beetles Green Cloverworm Green Peach Aphid ¹ Hornworms Meadow Spittlebug Pea Aphid ¹ Potato Leafhopper Spotted Alfalfa Aphid ¹ Threecornered Alfalfa Hopper Velvetbean Caterpillar Webworms	6.4 fl. oz/A (0.1 lb ai/A)	Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage. Use higher specified rate for increased pest pressure or for increased residual pest control. Do not exceed maximum allowable rate. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. Higher volumes of finished spray may improve insect control under higher temperatures, when foliage is dense and/or when insect pressure is high.
Armyworm, southern Armyworm, true Armyworm, yellowstriped Ant Cereal Leaf Beetle Chinch Bug Cricket Grass Mealybug Grasshoppers Range Caterpillar Stink Bugs Armyworm, fall Black Grass Bug		Do not apply more than 0.2 lbs. active ingredient per acre per season. Do not make applications less than 14 days Applications may be made up to 30 days prior to harvest for forage and hay.
Hunting Bill Bug Plant Bug spp.		¹ Aphid control may be variable depending on species present and host-plant relationships.

Read the Limitation of Warranty and Liability on the container labeling before buying or using BIFENTURE EC. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the container label.

Rev. 05/15/2015

United Phosphorus, Inc.
 630 Freedom Business Center, Suite 402
 King of Prussia, PA 19406
 1-800-438-6071