

Supplemental Label

BIFENTURE® EC Agricultural Insecticide

(EPA Reg. No. 70506-57)

DIRECTIONS FOR USE ON DRIED BEANS AND PEAS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application.

PESTS CONTROLLED	BIFENTURE EC/ACRE	APPLICATION INSTRUCTIONS	(days)
Flea Beetle Aster Leafhopper Leafhoppers Aphids Armyworms	0.025-0.10 lb ai (1.6-6.4 fl. oz. product) 0.033-0.10 lb ai (2.1-6.4 fl. oz.	Apply in a minimum of 10 gallons/A by ground or a minimum of 2 gallons/A by air. When applying by air, substitute 1-2 quarts of emulsified oil for 1-2 quarts water if desired. For best control thorough coverage is necessary.	14
Beet Fall Southern Yellowstriped Beetles Bean Leaf Cucumber Japanese (adult) Mexican Bean	product)	Do not apply more than 0.2 lb. a.i. (12.8 fl. oz. product) per acre per season to peas. Do not apply more than 0.3 lb. a.i. (12.8 fl. oz. product) per acre per season to beans. Do not make applications less than 7 days apart.	
Sap Bugs Plant Bugs Stink Bugs Tarnished Plant Bugs Alfalfa Caterpillar Cloverworm European Corn Borer			
Cutworms Western Bean Cutworm Corn Earworm Loopers (cont.)			
	Aster Leafhopper Leafhoppers Aphids Armyworms Beet Fall Southern Yellowstriped Beetles Bean Leaf Cucumber Japanese (adult) Mexican Bean Sap Bugs Plant Bugs Stink Bugs Tarnished Plant Bugs Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Loopers	Flea Beetle Aster Leafhopper Leafhoppers Aphids Armyworms Beet Fall Southern Yellowstriped Beetles Bean Leaf Cucumber Japanese (adult) Mexican Bean Sap Bugs Plant Bugs Stink Bugs Tarnished Plant Bugs Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Loopers	Flea Beetle Aster Leafhopper Leafhoppers Aphids Armyworms Beet Southern Yellowstriped Beetles Bean Leaf Cucumber Japanese (adult) Mexican Bean Sap Blugs Plant Bugs Stink Bugs Tarnished Plant Bugs Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Loopers Aphids (1.6-6.4 fl. oz. product) (2.1-6.4 fl. oz. product) Apply in a minimum of 10 gallons/A by ground or a minimum of 2 gallons/A by air. When applying by air, substitute 1-2 quarts of emulsified oil for 1-2 quarts water if desired. For best control thorough coverage is necessary. Do not apply more than 0.2 lb. a.i. (12.8 fl. oz. product) per acre per season to peans. Do not make applications less than 7 days apart.

Corn Rootworm (Adult) Thrips Webworms Pea Weevil Pea Leaf Weevil Whitefly Imported Cabbageworm Saltmarsh Caterpillar Tobacco Budworm Leafminers Grasshoppers		
Banks Grass Mite Carmine Mite Lygus Spp. Twospotted Spider Mite	0.08-0.10 lb ai (5.12-6.4 fl. oz. product)	

Rev. 5/09 CA appr. 10/28/09

> United Phosphorus, Inc. 630 Freedom Business Center King of Prussia, PA 19406 1-800-438-6071 • www.upi-usa.com