

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Dimension[®] 2EW

EPA Reg. No. 62719-542

For Distribution and Use Only in AL, AR, AZ, FL, GA, LA, MS, NC, NM, NV, OK, SC, TN, TX

Postemergent Crabgrass Control in Warm Season Turfgrass

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dimension[®] 2EW specialty herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dimension 2EW according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dimension 2EW.

Directions for Use

Refer to product label for Precautions and Restrictions, Mixing Instructions and Application Directions.

Use sites would include, established lawns, commercial sod farms, ornamental and sports turf (including sport fields, golf course fairways, roughs, tee boxes, unimproved turfgrass areas).

Postemergence Crabgrass Control in Warm Season Turf

This product provides both preemergence and postemergence control of crabgrass (including large, smooth, and southern species) in established lawns and ornamental turf. This product provides postemergence control of crabgrass through the 3 to 5 tiller stage of growth dependent upon location. The addition of a nonionic surfactant at a minimum of 0.25% v/v (2 pt per 100 gallons of spray) is recommended to improve postemergence control past the 5 leaf stage of growth. Read and follow the surfactant manufacturer's label recommendations. Postemergence control of this product can be improved by mowing turfgrass no sooner than two days prior and two days after application.

When applied at 2 pints per acre this product has demonstrated postemergent crabgrass control through the 3 to 5 tiller stage of growth in the western, southern and transition regions where warm-season turfgrasses are the predominate species.

For preemergence residual control of crabgrass, apply at least 0.5 inch of water after application; but in order to optimize postemergence control delay irrigation for 6 hours after application.

Use Restrictions:

- Do not apply this product to golf course putting greens.
- Do not harvest sod until 3 months or longer after application.
- Do not apply this product until the turfgrass has recovered from cultural practices such as verticutting or core-, spike-, or hydro-aerification.

- Do not use clippings from treated turf for mulching around vegetables or fruit trees.
- Do not apply this product through any type of irrigation system.
- Do not apply more than 2 pints of Dimension 2EW per acre (0.73 fl oz per 1000 sq ft) per application.
- Do not apply more than 6 pints of Dimension 2EW per acre per year (2.2 fl oz per 1000 sq ft).

Expiration Date: 6/30/2018

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
R337-004
EPA accepted 08/11/15
Initial printing