

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Treflan* TR-10*

EPA Reg. No. 62719-131

EPA 24(c) Special Local Need Registration SLN KS-990006
For Distribution and Use Only in the State of Kansas

Improved Control of Pigweed Species (Including Palmer Amaranth and Tall Waterhemp) and Dodder Weed in Established Alfalfa

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Treflan* TR-10* herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Treflan TR-10 according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Treflan TR-10.

Directions for Use

Location: Statewide

Crop/Site/Commodity: Established alfalfa

Target Pest/Problem: Pigweed species (including Palmer amaranth and tall waterhemp) and dodder weed. See "Alfalfa" section of product label for Treflan TR-10 for additional weeds controlled.

Dosage and Application Timing: Apply 20 lb Treflan TR-10 (2 lb a.i.) per acre for all soil textures prior to weed emergence. An additional 10 lb (1 lb a.i.) of Treflan TR-10 per acre may be applied approximately 60 days after the first application. Treflan TR-10 may be applied during dormancy or throughout the growing season immediately after a cutting. Because Treflan TR-10 does not control established weeds, application must be made prior to the expected time of weed germination.

Method of Application: Apply with properly calibrated ground or aerial broadcast equipment.

Activation: A single rainfall or overhead sprinkler irrigation of 0.5 inches or more, flood irrigation or furrow irrigation after application is required to activate Treflan TR-10. If activated using furrow irrigation, the surface of beds between furrows should be thoroughly wetted. If activation does not occur within 3 days after application, Treflan TR-10 should be activated using incorporation equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

Field Reentry After Application: See "Agricultural Use Requirements" section of product label.

Precautions and Restrictions:

- Do not cut or graze alfalfa within 21 days after application of Treflan TR-10.
- In the growing season following application of 20 or more pounds of Treflan TR-10 to alfalfa, plant only those crops for which Treflan TR-10 is registered as a preplant treatment or crop injury may occur.

*Trademark of Dow AgroSciences LLC
D06-048-003
Accepted: 05-14-99
Initial Printing