

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban* 15G

(EPA Reg. No. 62719-34)

2(ee) Recommendation [†]

For Distribution and Use Only in the State of Virginia

Use Recommendation for Control of Potato Leafhopper in Peanuts

ATTENTION

- [†] This recommendation is permitted under FIFRA 2(ee) and has not been submitted to or approved by the EPA or state lead pesticide agency.
- Read and follow all applicable directions for use, precautions and limitations on the product label for Lorsban* 15G insecticide.

Directions for Use

Lorsban 15G applied for soil insect control, such as southern corn rootworm, applied to peanuts at early flowering to peggings stage of growth in a 10-18" band over the row, will also provide control of potato leafhopper.

Restrictions

- Do not apply more than 15 ounces of Lorsban 15G per application per 1,000 feet of row.
- Do not apply more than 30 oz per 1000 ft of row or 26.6 pounds per acre per crop season.
- Do not harvest within 21 days after treatment.
- Do not feed peanut forage or hay to meat or dairy animals.

The combined total of preplant and postplant applications of Lorsban-4E and Lorsban 15G must not exceed 4 pounds of active ingredient per acre per crop season.

*Trademark of Dow AgroSciences LLC

D06-061-003

Issued: 04-23-02

Initial printing.