

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Lorsban\* 15G

EPA Reg. No. 62719-34

### Use of Lorsban 15G Prior to Post-Application of Lumax or Camix to Field Corn

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban\* 15G granular insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban 15G according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban 15G.

#### Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Soil applied Lorsban 15G, as a T-band or in-furrow application, may be followed by post-applied Lumax or Camix herbicide, applied in accordance with their label recommendations. Apply Lorsban 15G at-plant at a rate of 8 to 12 oz per 1000 feet of row to control soil insects such as corn rootworm, cutworm, wireworm, white grub, seed corn maggot, and others listed on the product label.

**Do not** use Lorsban 15G prior to the use of Lumax or Camix for corn grown for seed production due to the unknown nature of the response of inbred lines to Lumax or Camix.

\*Trademark of Dow AgroSciences LLC

D06-061-008  
EPA accepted: 06/03/03  
Initial printing.