

FIFRA SECTION 24(c) SUPPLEMENTAL LABELING
FOR DISTRIBUTION AND USE ONLY WITHIN TEXAS



Terramaster® 4EC

FUNGICIDE

FOR USE ON GREENHOUSE TOMATOES

EPA REG. NO. 400-422

EPA SLN NO. TX-030005

Active Ingredient: (% by weight)

Etridiazole [5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole]* 44.3%

Inert Ingredients:** 55.7%

Total: 100.0%

*Contains 4 lbs. of 5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole per U.S. Gallon at 20°C.

**Contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN DANGER • PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of State and Federal law to use this product in a manner inconsistent with its labeling.

GREENHOUSE TOMATOES

For control of *Pythium* and *Phytophthora* Root Rot in
greenhouse tomatoes by drip irrigation

Drip Irrigation

Apply 6 - 7 fluid ounces of TERRAMASTER 4EC per acre in a 0.01% solution (6.5 fluid oz. in 500 gallons of water) no sooner than 3 weeks after transplanting.

Reapply as needed, 6 - 7 fluid ounces of TERRAMASTER 4EC per acre in a 0.01% solution, no sooner than 3 weeks after a previous application.

Do not exceed 27.4 fluid ounces of TERRAMASTER 4EC per acre per crop season.

Do not apply solutions more concentrated than 0.01% without first verifying potential phytotoxicity on the variety of tomato being grown.

Do not apply TERRAMASTER 4EC sooner than 3 days before harvest.

Uniform distribution of TERRAMASTER 4EC through the growing medium is essential for best results. Localized over-dosing may cause crop damage and under-dosing may cause ineffective disease control.

TERRAMASTER 4EC should be introduced into the growing medium only after the medium has been thoroughly irrigated to ensure adequate product movement through the growing medium. Continue irrigation until the entire desired quantity of TERRAMASTER 4EC per acre has been introduced.

Do not mix TERRAMASTER 4EC with other pesticides or fertilizers unless compatibility tests have shown complete compatibility.

Apply TERRAMASTER 4EC to a small crop area to ensure crop safety before applying to broad crop areas.

THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER

Follow all applicable directions, restrictions, and precautions on the EPA registered label.

This label must be in the possession of the user at the time of the pesticide application.

002

Terramaster® is a Registered Trademark of Chemtura Corporation

24c (SLN) REGISTRANT:

Chemtura Corporation

199 Benson Road

Middlebury, CT 06749

EPA SLN NO. TX-030005