Supplemental Label



ESTEEM® ANT BAIT Insect Growth Regulator EPA Reg. No. 59639-114

ESTEEM ANT BAIT INSECT GROWTH REGULATOR USES ON: BUSHBERRIES, FIELD CROP & VEGETABLES, TREE FRUIT & NUTS AND TROPICAL & SUB-TROPICAL FRUITS

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

CROPS	PESTS	RATES PER ACRE	SPECIAL INSTRUCTIONS
Bushberries:	Ants	1.5 to 2.0 lbs.	Broadcast or Aerial Application: Apply
Blueberry	Pavement Ant		uniformly with ground or aerial equipment
Currant	Red Imported Fire Ant		calibrated to give the correct dosage.
Elderberry	Southern (California) Fire		
Gooseberry	Ant		ESTEEM Ant Bait should be applied in the
Huckleberry			early Spring or Summer at the first sign of
Juneberry			fire ant activity. Applications may be made
Lingonberry			any time of the day but is more effective
Salal			when ants are actively foraging, usually
Field Crops &			when the soil temperature is above 60° F.
Vegetables:			Avoid application if rain is expected within
Cotton			4 to 6 hours. Do not water treated area for
Eggplant			24 hours after application. In the event of
Ground Cherry			prolonged rains after initial treatment,
Okra			reapplication may be necessary.
Pepino			
Pepper			A second application of ESTEEM Ant Bait
Tomatillo			may be advisable after 12 to 16 weeks in
Tomato			areas of heavy infestations, or when the
Tree Fruits & Nuts:			mounds remain active, or when a
Almond			reinfestation occurs. In large mounds,
Apple			some worker fire ants may persist for a
Apricot			while after the queen is dead. For 7 to 10
Cherry, Sweet & Tart			days after treatment, do not apply any
Citrus			other fire ant pesticide. This is to allow
Fig			sufficient time for the worker ants to seek
Nectarine			out the bait and take it back to the mound.
Nut Trees			After 14 days, it may be desirable to apply
Olive			a registered fire ant pesticide to kill
Peach			remaining foraging ants.
Pear			Design Almond Timbre 1
Pistachio			Bearing Almond Timing: In bearing
Plum			almond orchards apply at 6 to 8 weeks
Walnut			prior to harvest to protect fallen nuts.

Continued

CROPS, Continued	PESTS	RATES PER ACRE	SPECIAL INTRUCTIONS
Tropical & Subtropical	Ants	1.5 to 2.0 lbs.	Broadcast or Aerial Application: Apply uniformly
Fruits:	Pavement Ant		with ground or aerial equipment calibrated to give the
Acerola	Red Imported Fire Ant		correct dosage.
Atemoya	Southern (California)		
Avocado	Fire Ant		ESTEEM Ant Bait should be applied in the early
Biriba			Spring or Summer at the first sign of fire ant activity.
Black Sapote			Applications may be made any time of the day but is
Canistel			more effective when ants are actively foraging,
Cherimoya			usually when the soil temperature is above 60° F.
Custard Apple			Avoid application if rain is expected within 4 to 6
Feijoa			hours. Do not water treated area for 24 hours after
Guava			application. In the event of prolonged rains after
llama			initial treatment, reapplication may be necessary.
Jaboticaba			
Longan			A second application of ESTEEM Ant Bait may be
Lychee			advisable after 12 to 16 weeks in areas of heavy
Mamey Sapote			infestations, or when the mounds remain active, or
Mango			when a reinfestation occurs. In large mounds, some
Passionfruit			worker fire ants may persist for a while after the
Papaya			queen is dead. For 7 to 10 days after treatment, do
Pulasan			not apply any other fire ant pesticide. This is to allow
Rambutan			sufficient time for the worker ants to seek out the bait
Sapodilla			and take it back to the mound. After 14 days, it may
Soursop			be desirable to apply a registered fire ant pesticide to
Spanish Lime			kill remaining foraging ants.
Star Apple			<u> </u>
Starfruit			Bearing Almond Timing: In bearing almond
Sugar Apple			orchards apply at 6 to 8 weeks prior to harvest to
Wax Jambu			protect fallen nuts.

Individual Mound Treatment: Apply 2 to 4 level Tbsps. of bait per mound, uniformly distributing material 3 to 4 ft. around the mound. Do not disturb the mound. Do not apply to top of mounds.

- · Do not allow livestock to graze treated areas.
- · Do not apply within 24 hours of harvest.

MODE OF ACTION: ESTEEM Ant Bait is an Insect Growth Regulator similar to the naturally occurring insect growth hormones which interferes with (stops) their growth or development. ESTEEM Ant Bait breaks the reproductive life cycle of ants, causing starvation of the colony. Ant workers pick up the bait granules and take them back to the colony. Workers feed the bait to both the queen and immature ants thereby preventing worker replacement. Due to degeneration of reproductive organs, the ant queen cannot replace workers. The lack of worker replacement results in colony death as the existing worker ants age and die. Within 3 to 4 weeks there is substantial colony mortality and within eight weeks the majority of the colony population has been eliminated.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. PLEASE REFER TO CONTAINER LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

PLEASE CONTACT VALENT U.S.A. CORPORATION AT 1-800-6-VALENT (682-5368) TO DETERMINE IF THIS USE IS REGISTERED IN YOUR STATE.

Copyright © 2003, 2004 by Valent U.S.A. Corporation ESTEEM - Reg. TM of Valent U.S.A. Corporation

Manufactured for Valent U.S.A. Corporation | P. O. Box 8025 | Walnut Creek, CA 94596-8025 | www.valent.com

Made in U.S.A.

Form: 2003-EAB-0010



Products That Work, From People Who Care™ | www.valent.com | 800-6-VALENT (682-5368)

2003-EAB-0010 5/04 (mf)