Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lock-On[®]
EPA Reg. No. 62719-79

EPA 24(c) Special Local Need Registration SLN TX-040027 (For Distribution and Use Only in the State of Texas)

For Use in Pink Bollworm Control in Cotton

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lock-On[®] insecticide before applying. All applicable directions, restrictions and precautions on EPA registered product labeling must be followed.
- Use of Lock-On according to this supplemental labeling is subject to all use precautions and limitations imposed
 by the label affixed to the container for Lock-On.

Directions for Use

Apply as a broadcast foliar spray using aircraft or ground spray equipment. Use sufficient spray volume to ensure thorough coverage of treated plants, but no more than 10 gallons per acre (gpa) using aerial (fixed wing or helicopter) or power operated ground equipment. Do not exceed 10 gpa total spray volume. Increase spray volume when foliage is dense and/or pest population is high and/or under high temperature and wind conditions. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control.

| | Application Rate |
|---------------|------------------|
| Pest | (pint/acre) |
| pink bollworm | 2 |

Specific Use Restrictions:

- Preharvest Interval: Do not apply within 40 days before harvest.
- Do not make more than 6 applications of Lock-On or other chlorpyrifos containing product per crop season or apply more than 3 lb a.i. chlorpyrifos per acre per season.
- Do not make a second application of Lock-On or other product containing chlorpyrifos within 10 days of the first application.
- Do not allow livestock to graze in treated areas.
- Do not feed gin trash or treated forage to livestock.
- Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours unless PPE required for early entry is worn.
- Chemigation: Do not apply Lock-On through any type of irrigation system.

24(c) Registrant: Dow AgroSciences LLC 9330 Zionsville Rd Indianapolis, IN 46268

[®]Trademark of Dow AgroSciences LLC R060-002 Approved: 06/23/04 Initial printing.