

# GC-3

**Fungicide For Powdery Mildew Control**  
**Controls Powdery Mildew on Agricultural**  
**Crops**



JH Biotech, Inc.  
 Ventura, Ca  
 TEL: (800) 428-3493  
 www.jhbiotech.com

This product is exempt from registration with the Federal EPA under section 25(b) of FIFRA. GC-3 has not been registered with the Environmental Protection Agency. JH Biotech, Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act.

**U.S. Patented and Patent Pending**

## GC-3 APPLICATION & USE TABLE

GC-3 is a contact fungicide.  
 GC-3 is exempt from tolerances on all food crops.

CROPS	DISEASE	RATE PER ACRE	
Beans & Peas Dry Beans Fresh Beans Berries Strawber- ries Raspberries Fruit& Nut Trees Pome Fruit Stone Fruit Grapes Leafy Vegetables Lettuce Spinach Cole Crops Melons & Cucurbits Muskmelon Watermelon Cucumber Squash	Powdery Mildew	Mix 1 gallon of GC-3 concentrate per 100 gallons of spray water. Apply 100-200 gallons of solution per acre, or enough to cover the entire surface of infected tissues.  <u>Mix Volume</u> 1 Gallon 5 Gallon 10 Gallons 25 Gallons 100 Gallons	<u>FL. OZ.</u> 1.5 7 14 32 128
		Apply at intervals of 7-10 days. Repeat application as necessary. Coverage is essential to establish control. The use of a spreader/ sticker may increase contact and efficacy of treatment.	

## MIXING INSTRUCTIONS

GC-3 may be applied using conventional ground application equipment. Aerial application equipment is not recommended. Use quantities of water sufficient to provide thorough coverage of infested plants. Add enough water to spray tank to allow maximum agitation. With agitator running, slowly add the appropriate amount of GC-3. Continue agitation. Add remainder of water and agitate until thoroughly mixed. Maintain suspension while loading and spraying. Do not mix more GC-3 than can be used in a 12 hour period. Rinse and flush spray equipment thoroughly following each use. Do not contaminate water when disposing of equipment washwaters.

In order to make proper decisions on application rates to be used, follow the recommendations in the **APPLICATION RATE TABLE**.

**NOTE: All applications should be proceeded by a phytotoxicity check to ensure that the material is safe for that particular plant variety.** We recommend that prior to treatment of a large number of plants, new species, or varieties, treat a few plants with GC-3 and observe if there is any phytotoxicity. Certain varieties of peppers are sensitive to GC-3. Treat a small number of plants with GC-3 before treating the whole area. It is best to treat plants after they've adjusted to environmental changes, i.e. temperature, water, light intensity, new containers, new area, transplanting, wind, etc.

## TANK MIXING

GC-3 has not been tested for use in a tank mix with many commonly used insecticides, fungicides, or spray tank adjuvants. We recommend testing all tank mix combinations for physical compatibility and plant safety. Consult your dealer or manufacturer for specific recommendations. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rate should be exceeded. Application must be made in accordance with the more restrictive of label limitations and precautions. Read and follow label instructions.

\* May cause phytotoxicity to peppers. See Note.