



## SUPPLEMENTAL LABELING

**RESTRICTED USE PESTICIDE  
DUE TO AVIAN AND AQUATIC TOXICITY  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT  
SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION**

### DIAZINON 50W

A Wettable Powder Insecticide  
EPA REG NO. 66222-10

**FOR USE ON FIGS IN CALIFORNIA, WATERCRESS IN HAWAII, AND FILBERTS IN OREGON AND  
WASHINGTON**

ACTIVE INGREDIENT:	% BY WT.
Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate .....	50%
INERT INGREDIENTS: .....	50%
	<b>TOTAL 100%</b>

## KEEP OUT OF REACH OF CHILDREN CAUTION

#### DIRECTIONS FOR USE

It is violation of federal law to use this product in a manner inconsistent with its labeling.  
This labeling must be in the possession of the user at the time of pesticide application.  
All applicable directions, restrictions, and precautions on the EPA-registered label are to be followed.

Crop	Pest	Rate	Comments
<b>FIGS</b> (California Only)	Vinegar flies- <i>drosophila</i> spp., Dried fruit beetles	Apply 1 lb. per 100 gals of water not to exceed 200 gals. of water per application	Apply as infestations occur. Repeat application after 7 days if necessary.
	<ul style="list-style-type: none"> <li>Do not apply more than 2 pounds of product per acre per application and no more than 6 pounds of product per acre per year.</li> <li>To avoid illegal residues, allow a minimum of 5 days between the last application and harvest.</li> </ul>		
<b>FILBERTS (HAZELNUTS)</b> (Oregon and Washington Only)	Filbert Leafrollers, Aphids	Apply 2-4 lbs. in 250-400 gals. of water per acre	Apply as a thorough cover spray. Repeat application after 14 days if necessary.
	<ul style="list-style-type: none"> <li>Do not apply more than 4 pounds of product per acre per application and no more than 12 pounds per year.</li> <li>To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.</li> </ul>		

<b>Crop</b>	<b>Pest</b>	<b>Rate</b>	<b>Comments</b>
<b>WATERCRESS</b> (Hawaii Only)	Cyclamen Mites	Apply 1 pound per 100 gals. of water per acre.	Apply as a thorough cover spray. Spray should be directed into the crowns and growing tips of plants. If infestation is heavy, make a second application within 10 days after the first application.
	<ul style="list-style-type: none"> <li>• Do not apply more than 1 pound of product per acre per application and no more than 2 pounds of product per acre per season.</li> <li>• To avoid illegal residues; allow a minimum of 5 days between the last application and harvest.</li> </ul>		

Diazinon 50W (66222-10)(supplemental on figs, filberts, watercress EPA app 5-15-02)(CA app 11-23-04)