

For Agricultural or Commercial Use Only EPA REG. No. 279-3158

EMERGENCY CALLS: 800-331-3148



FOR DISTRIBUTION AND USE ONLY IN TEXAS

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH IT LABELING

Section 24(c) State Registration

DIRECTIONS FOR USE:

For aerial application of Command 3ME on rice. No diluted spray of Command 3ME herbicide shall be applied by air in the following counties: Harris, Fort Bend (North and East of Highway 36).

Apply prior to weed emergence during the period from 14 days pre-plant to 7 days post - plant or apply post-plant to rice at 1-2 leaf stage to provide pre-emergent and extended residual grass control. If grass is already present, include a post emergence grass herbicide with the application. Command 3ME may only be tank-mixed with one tank mix partner among the following herbicides: AIM, Bolero, Buccaneer Plus, Clincher, Cornerstone, Credit Extra, Duet, Durango, Facet, Glyphomax Plus, Glyfos Xtra, Grasp, Makaze, Mirage Plus, Newpath, Permit, QuinStar, QuinStar 4L, Rattler, Ricestar, Stam, Super Wham, as well as Crop Oil Concentrate or Non-ionic surfactant. Only a single application of Command 3ME plus an approved glyphosate formulation is allowed per season.

The following herbicides may be added to Command 3ME + AIM for a three way tank mix: Facet, Newpath, Permit, Ricestar, or Super Wham. The following herbicides may be added to Command 3ME + Grasp SC for a three / four way tank mix: Facet, Facet + Permit, Grandstand R, Permit, Quinstar, Quinstar + Permit, or Super Wham. The following three way tank mix of Command + Super Wham / Stam + Permit may be used. Crop Oil Concentrate or Non-ionic surfactant may be applied with Command 3 ME alone or with tank-mixes listed above.

Observe all buffer restrictions noted in the Application Precautions and Restrictions sections.

Partial weed control may result if levees are formed after Command 3ME has been applied.

Additional use of labeled post-emerge herbicide applications may be required where existing grass weeds are present at the time of application.

CROP	PESTS	RATE OF APPLICATION			
Rice	Barnyardgrass (Watergrass) (Including herbicide resistant types) Broadleaf Signalgrass Crabgrass (Large, Smooth)	Soil Texture	Broadcast Rates Per Acre*	Pounds active ingredient per acre	Acres per Gallon
	Panicum (Common, Fall, Texas) Sprangletop	Coarse (light) Soils: (sand,loamy sand, sandy loam):	0.67-0.8 pts (10.7 - 12.8 fl oz)	0.25 - 0.3	12.0 - 10.0
		Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1.1 pts (17.1 fl oz)	0.4	7.5
		Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1.33 - 1.6 pts (21.3 - 25.6 fl oz)	0.5 - 0.6	6 - 5
		* Select lower to higher rates based on lighter to heavier soil types.			



24(C) Registrant: FMC Corporation Agriculture Products Group 1735 Market Street Philadelphia, PA 19103

CROP SAFETY PRECAUTIONS:

Application of Command 3ME herbicide to fields that have been precision leveled with deep cuts may result in rice crop injury including stand loss. Consult with rice specialists for soil amending practices that can reduce potential for herbicide injury in precision-leveled fields.

AERIAL APPLICATION REQUIREMENTS:

Aircraft used to apply Command 3ME shall be configured and operated in such manner as to minimize off-site spray movement to desirable species.

APPLICATION PRECAUTIONS:

Caution must be taken to minimize off target drift of Command 3ME as off-site movement of Command can cause foliar whitening or yellowing of some plants.

Do not apply Command 3ME herbicide within 1,200 feet of the areas listed below:

Towns and Housing Developments

Commercial Vegetable Production (except peppers, pumpkins, succulent peas, sweet corn, sweet potato and winter squash)

Commercial Fruit / Nut Production

Commercial Nurseries

Commercial Greenhouses

Avoid making applications of Command 3ME herbicide when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Prior to application, adjacent properties must be checked, and applications within 300 feet of desirable plants must be avoided.

RESTRICTIONS:

Command 3ME may be applied to water seeded rice 14 days prior to planting or during pegging up to re-flooding but prior to grass emergence.

With split applications do not apply more than a total of 1.6 pt/A Command 3ME (0.6 lbs ai/A) per season.

Do not apply Command 3ME herbicide on rice fields in which concurrent crayfish or catfish farming is included in the cultural practices. Do not use water containing Command 3ME residues from rice cultivation to irrigate food or feed crops, which are not registered for use with Command 3ME.

No diluted spray of Command 3ME herbicide shall be applied by air in the following counties: Harris, Fort Bend (North and East of Highway 36

"This product should only be applied when the potential for drift to adjacent sensitive areas (known habitat for endangered species) is minimal (wind is blowing away from sensitive area)".

REPLANTING INSTRUCTIONS

If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with Command 3ME herbicide. Do not retreat fields with a second application of Command 3ME. Do not replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP GUIDELINES on this labeling.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

Note: When using Command 3ME herbicide with other registered herbicides always refer to rotational restrictions and precautions on the other product s label.

ANYTIME - Cotton*, Peas, Peppers, Pumpkins, Rice, Soybeans, Tobacco

9 MONTHS - Cotton, Corn (Field, Pop, Seed, Sweet), Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Sweet Potatoes, Tomatoes (Transplanted)

12 MONTHS - All crops

 Refer to "REQUIREMENTS FOR PLANTING TIME APPLICATIONS" and "REPLANTING INSTRUCTIONS" in the Command 3ME Section 3 label.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.



24(C) Registrant: FMC Corporation Agriculture Products Group 1735 Market Street Philadelphia, PA 19103