

# FIFRA Section 2(ee) Recommendation



FOR DISTRIBUTION AND USE ONLY WITHIN THE FOLLOWING COUNTIES IN THE STATE OF TEXAS: ATASCOSA, BASTROP, BEXAR, BOSQUE, BROWN, COMANCHE, DENTON, DE WITT, DUVAL, EASTLAND, ERATH, FAYETTE, FRIO, GRAYSON, HARRIS, HIDALGO, HOOD, HOUSTON, LAMPASAS, LA SALLE, LAVACA, LEE, MASON, MC CULLOCH, MEDINA, PARKER, ROBERTSON, STEPHENS, WALKER, WILSON



FOR POSTEMERGENCE WEED CONTROL IN PEANUTS  
EPA Reg. No. 59639-34  
FIFRA SECTION 2(ee) RECOMMENDATION

## DIRECTIONS FOR USE

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

**USE COBRA® HERBICIDE AT 12.5 OZ/A FOR POSTEMERGENCE CONTROL OF THE FOLLOWING WEEDS IN PEANUTS.**

### *Cobra* Herbicide Peanuts – Rate, Weeds and Weed Sizes

Product	Product Rate	Broadleaf Weeds Controlled	Maximum Weed Size
<i>Cobra</i> Herbicide + Crop Oil Concentrate	12.5 oz/A + Refer to specific Information on the next page	Palmer Amaranth Smellmelon Horse Purslane	6 inches 8 inches 8 inches

Refer to the *Cobra* Herbicide label for a complete list of weeds controlled.

Do not apply *Cobra* Herbicide until peanuts have at least six true leaves.

After at least 14 days, a second application of *Cobra* Herbicide at 12.5 oz/A may be made to control later emerging weeds or weeds escaping the first application.

*Cobra* Herbicide may be applied sequentially or as a tank mix with other herbicides for additional grass and broadleaf weed control. Refer to the product label for specific recommendations.

### Use Restrictions:

1. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.
2. Do not apply more than 12.5 fl oz/A of *Cobra* Herbicide per application.
3. Do not apply more than 25.0 oz of *Cobra* Herbicide per acre during a single growing season.
4. Do not graze treated fields or feed treated forage or hay to livestock.
5. Do not apply *Cobra* Herbicide to peanuts later than 90 days before harvest.
6. Do not apply this product through any type of irrigation equipment.

## APPLICATION INFORMATION

**Ground Application:** Apply *Cobra* Herbicide on a broadcast basis with 10 to 20 gals of water per acre and a spray pressure of 40 to 60 psi. Use flat fan or hollow cone nozzles and adequate spray volume to ensure thorough weed coverage. Dense leaf canopies may shield smaller weeds and prevent adequate coverage. If crop or weed foliage is dense, increase water volume, up to 50 gals/A.

When row banding, best coverage is provided when two nozzles are directed from either side of the row towards the weeds in the middle. The recommended minimum band width is 15 inches, with a minimum spray volume of 15 gpa on the band. Band application using a single nozzle over the row is not recommended.

Adjuvant Recommendations: Use COC at 1 to 2 pts/A. Under less than ideal growing conditions or when targeting weeds near the top of the height range, use the higher COC rate. Only under ideal growing conditions and when targeting weeds at the lower end of the height range (refer to product label), should non-ionic surfactant be used at 0.25% v/v (1 quart per 100 gals of water).

**Aerial Application:** *Cobra* Herbicide may be applied by air using 5 to 10 gals of water per acre. Applications at less than 5 gpa will provide inadequate control. Use precautions on the product label to minimize drift.

Adjuvant Recommendations: Use crop oil concentrate at the rate of 1% v/v in the spray solution. In dry conditions, use at least 2% v/v of crop oil concentrate.

\*\*\*\*\*

Follow all applicable directions, restrictions and precautions on the EPA registered labels.

It is a violation of Federal law to use these products in a manner inconsistent with their labeling.

Registrant: Valent U.S.A. Corporation  
P.O. Box 8025  
Walnut Creek, CA 94596-8025

*Cobra* is a registered trademark of Valent U.S.A. Corporation