

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Stinger[®]

EPA Reg. No. 62719-73

24(c) Special Local Need Registration EPA SLN No. NC-030005

Postemergence Broadleaf Weed Control in Strawberry
(For Distribution and Use Only in the State of North Carolina)

Special Conditions and Risks of Use

USE OF Stinger[®] herbicide (THE "PRODUCT") ON STRAWBERRY (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Dow AgroSciences makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;
- (3) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Special Conditions and Risks of Use supersede any contrary representations or recommendations by Dow AgroSciences, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Dow AgroSciences and not specified by the U.S. EPA or the State of North Carolina.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Stinger® herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Stinger is subject to all use precautions and limitations imposed by the label affixed to the container for Stinger.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Note: Some leaf cupping may occur. Do not use if unwilling to accept leaf injury and possible yield reduction.

Perennial (MATTED ROW) Strawberry Production

Target Broadleaf Weeds, Application Rate, and Use Restrictions:

Target Broadleaf Weeds	Stinger (pint/acre)	Use Restrictions
artichoke, Jerusalem clovers cocklebur, common curly dock dandelion groundsel jimsonweed lettuce, prickly nightshade ragweed, common ragweed, giant smartweed (suppression) sorrel, red sowthistle, annual (suppression) sunflower thistle, bull (suppression) thistle, Canada (suppression) vetch	Spring application: 1/3 Post-harvest application: 1/3 - 2/3	<ul style="list-style-type: none"> • Make 1 to 2 applications per year not to exceed 2/3 pint per acre (0.25 lb ae per acre) per year. • Do not use Stinger with a surfactant on strawberries. • Do not tank mix with other pesticides. • Preharvest Interval: Do not apply within 30 days of harvest.

Broadcast Applications

Apply uniformly with ground equipment in 20 to 75 gallons of water per acre.

Spring Application: If needed, make a ground application in the spring, prior to bloom, no later than 30 days prior to harvest at 1/3 pint per acre. **Note:** If newly planted, **do not** make an application until strawberry plants are established; if needed on newly planted crop, make a post-harvest application of that year and, if needed, a spring application the following year.

Post-Harvest Application: Make a ground application after harvest at 1/3 to 2/3 pint per acre. For control of Canada thistle, apply Stinger after the majority of basal leaves have emerged but prior to bud stage of the weed.

Spot Applications: Refer to the Hand-Held Sprayers section of the label for Stinger for spot application instructions.

Recommendations:

- For difficult to control weeds, best results will be achieved with a post-harvest application followed by a spring application.

Annual (PLASTICULTURE) Strawberry Production

Target Broadleaf Weeds, Application Rate, and Use Restrictions:

Target Broadleaf Weeds	Stinger (pint/acre)	Use Restrictions
artichoke, Jerusalem clovers cocklebur, common curly dock dandelion groundsel jimsonweed lettuce, prickly nightshade ragweed, common ragweed, giant smartweed (suppression) sorrel, red sowthistle, annual (suppression) sunflower thistle, bull (suppression) thistle, Canada (suppression) vetch	<p>After weed emergence but at least 30 days PHI: 1/3 – 1/2</p> <p>Row middles: 1/3 - 2/3</p>	<ul style="list-style-type: none"> • Apply as a banded application over the strawberry row(s) or to the row middles, between plastic rows, or as a spot treatment. • Make 1 to 2 applications per year not to exceed 2/3 pint per acre (0.25 lb ae per acre) per year. • Do not apply Stinger until strawberries have established. • Do not use Stinger with a surfactant on strawberries. • Do not tank mix with other pesticides. • Preharvest Interval: Do not apply within 30 days of harvest. • Do not apply as a broadcast treatment.

Application

Apply to strawberries after weed emergence but at least 30 days prior to harvest. Use 1/3 – 1/2 pint per acre of Stinger post applied. Use up to 2/3 pint per acre for row middles.

Other Use Restrictions:

- **Do not** apply by aircraft
- **Chemigation:** Do not apply Stinger through any type of irrigation system.
- **Do not** apply within 6 to 8 hours of expected rainfall or irrigation.
- **Do not** compost treated vegetation if compost will be used on sensitive plants.
- Avoid injury to non-target plants. Refer to the product label.

®Trademark of Dow AgroSciences LLC

R043-023

File Code: NC-030005

Accepted: 06/21/06

Replaces D06-043-089.