

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

WideMatch™

EPA Reg. No. 62719-512

Conservation Reserve Program (CRP) Acreages - Permanent Grasses Only Non-Cropland: Including Fencerows, Farm Building Sites and Equipment Pathways

(For Distribution and Use In CO, KS, MI, MN, MT, ND, NE, OH, SD, WY)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for WideMatch™ herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of WideMatch according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for WideMatch.

Directions for Use

Application Timing: Apply as a broadcast postemergence treatment control of broadleaf weeds in established perennial grasses. Apply when weeds are actively growing, but before weeds are 8 inches tall or are vining. To obtain season-long control of perennial weeds such as Canada thistle, apply after the majority of the weed's basal leaves have emerged up to bud stage. Later applications may result in less consistent control.

Conditions of plant stress, such as drought, will increase potential for injury to grasses at all stages of growth. Do not apply to newly seeded areas grasses until well established. Perennial grasses are considered well established tillers and secondary roots have developed and growing vigorously.

Broadcast Application Rates:

(Numbers in parentheses (-) refer to footnotes following table.)

Weed Size or Species (1)	Application Rate (pt/acre)
Susceptible broadleaf weed seedlings less than 8 inches tall or vining (2)	1.33 – 2.66

1. See "Weeds Controlled or Suppressed" section for a complete listing of weeds controlled or suppressed.
2. Control of small kochia will be more consistent if kochia is at least 1 inch tall.

Tank Mixtures for Conservation Reserve Program (CRP) Acreages and Non-Cropland

WideMatch can also be tank mixed with 1/2 to 1 lb per acre of 2,4-D where target weeds are susceptible to 2,4-D. See "Tank Mixing Precautions" under "Mixing Instructions".

Precautions:

- **Rotation to Broadleaf Crops:** Do not plant broadleaf crops in treated areas until an adequately sensitive bioassay shows that no detectable clopyralid is present in the soil. (See Crop Rotation Intervals in the General Information section of the package label.)
- Do not use on CRP acreages or non-cropland that is underseeded with desirable legumes, clovers, or other sensitive broadleaf plants.

Restriction:

- Do not apply more than 5.33 pints or WideMatch per acre per use season on non-cropland areas or CRP acres.

™Trademark of Dow AgroSciences LLC

R202-002

Accepted 05/06/05

Initial printing.