

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Garlon* 3A

EPA Reg. No. 62719-37

Wetland Sites in Production Forests and Industrial Non-Crop Areas

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Garlon* 3A specialty herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Garlon 3A according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Garlon 3A.

Directions for Use

Garlon 3A may be used within production forests and industrial non-crop sites to control target vegetation in and around standing water sites, such as marshes, wetlands, and the banks of ponds and lakes and transition areas between upland and lowland sites.

For control of woody plants and broadleaf weeds in these sites, follow use directions and application methods on this label for forestry and terrestrial non-cropland sites.

Plants Controlled by Garlon 3A

Woody Plant Species

alder	Douglas-fir	poplar
arrowwood	dogwood	salt-bush (<i>Baccharis</i> spp.)
ash	elderberry	sassafras
aspen	elm	scotch broom
bear clover (bearmat)	gallberry	sumac
beech	hazel	sweetbay magnolia
birch	hornbeam	sweetgum
blackberry	kudzu†	sycamore
blackgum	locust	tanoak
Brazilian pepper	madrone	thimbleberry
cascara	maples	tulip poplar
ceanothus	mulberry	waxmyrtle
cherry	oaks	western hemlock
chinquapin	persimmon	wild rose
choke cherry	pine	willow
cottonwood	poison ivy	winged elm
crataegus (hawthorn)	poison oak	salmonberry

†For complete control, retreatment may be necessary.

Annual and Perennial Broadleaf Weeds

bindweed	dandelion	ragweed
burdock	field bindweed	smartweed
Canada thistle	lambsquarter	tansy ragwort
chicory	plantain	vetch
curly dock	Purple loosestrife	wild lettuce

Maximum Labeled Rate versus Spray Volume per Acre

Total Spray Volume (gal/acre)	Maximum Rate of Garlon 3A	
	Forestry Sites [†] (gal/100 gal of spray)	Other Non-Cropland Sites ^{††} (gal/100 gal of spray)
400	0.5	0.75
300	0.67	1
200	1	1.5
100	2	3
50	4	6
40	5	7.5
30	6.65	10
20	10	15
10	20	30

[†] Do not exceed the maximum use rate of 6 lb ae of triclopyr (2 gal of Garlon 3A)/acre/year.

^{††} Do not exceed the maximum use rate of 9 lb ae of triclopyr (3 gal of Garlon 3A)/acre/year on non-cropland use sites other than rangeland, pasture, forestry, and grazed areas.

Precautions for Potable Water Intakes for Emerged Aquatic Weed Control

See chart below for specific setback distances near functioning potable water intakes. **Note:** Existing potable water intakes which are no longer in use, such as those replaced by potable water wells or connections to a municipal water system, are not considered to be functioning potable water intakes. These setback restrictions do not apply to terrestrial applications made adjacent to potable water intakes.

Area Treated (acres)	Garlon 3A Application Rate, qt/acre			
	2 qt/acre	4 qt/acre	6 qt/acre	8 qt/acre
Setback Distance (ft)				
4	0	200	400	500
>4 - 8	0	200	700	900
>8 - 16	0	200	700	1000
>16	0	200	900	1300

To apply Garlon 3A around and within the distances noted above from a functioning potable water intake, the intake must be turned off until the triclopyr level in the intake water is determined to be 0.4 parts per million (ppm) or less by laboratory analysis or immunoassay.

- **Recreational Use of Water in Treatment Area:** There are no restrictions on use of water in the treatment area for recreational purposes, including swimming and fishing.
- **Livestock Use of Water from Treatment Area:** There are no restrictions on livestock consumption of water from the treatment area.

Use Precautions and Restrictions:

- **Do not** apply to salt water bays or estuaries.
- **Do not** apply directly to un-impounded rivers or streams.
- **Do not** apply on ditches or canals used to transport irrigation water. It is permissible to treat non-irrigation ditch banks.
- **Do not** apply where runoff water may flow onto agricultural land as injury to crops may result.
- When making applications to control unwanted plants on banks or shorelines of moving water sites, minimize overspray to open water.
- The use of a mistblower is not recommended.
- On forestry sites, Garlon 3A may be used at rates up to 6 lb ae of triclopyr (2 gallons of Garlon 3A) per

acre per year.

- For all terrestrial use sites other than range, pasture, forestry sites, and grazed areas, the maximum application rate is 9 lb ae of triclopyr (3 gallons of Garlon 3A) per acre per year.
- Minimize overspray to open water when treating target vegetation in and around non-flowing, quiescent or transient water. When making applications to control unwanted plants on banks or shorelines of flowing water, minimize overspray to open water.
- Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat such areas.

*Trademark of Dow AgroSciences LLC	
------------------------------------	--

D06-101-002 EPA-accepted 12/03/02 Initial printing.	
---	--