

# FIFRA Section 24(c) Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger<sup>®</sup>

EPA Reg. No. 62719-73

EPA SLN No. OR-030031

For Distribution and Use Only in the State of Oregon

This label is valid until December 31, 2019 or until otherwise amended, withdrawn, canceled, or suspended.

### Postemergence Broadleaf Weed Control in Strawberry

Use of Stinger<sup>®</sup> herbicide on strawberry may result in crop injury, crop yield reduction and/or crop loss. Read and understand the Special Conditions and Risks of Use section on this label before using this product in accordance with this FIFRA Section 24(c) Special Local Need.

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of pesticide application.
- Read this SLN and the label affixed to the container for Stinger<sup>®</sup> herbicide before applying.
- Use of Stinger according to this SLN is subject to all use directions, precautions and limitations imposed by the label affixed to the container for Stinger.

#### Active Ingredient:

clpyralid: 3,6-dichloro-2-  
pyridinecarboxylic acid,  
monoethanolamine salt.....40.9%

Other Ingredients .....59.1%

Total .....100.0%

Acid Equivalent: clpyralid: 3,6-dichloro-2-pyridinecarboxylic acid - 31% (3 lb/gal)

**Keep Out of Reach of Children**

## CAUTION

#### Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Do not contaminate water used for irrigation or domestic purposes.

Clopyralid is a chemical which can travel (seep or leach) through soil and under certain conditions contaminate groundwater which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an underlying aquifer is shallow, or to soils containing sinkholes over limestone bedrock, severely fractured surfaces, and substrates which would allow direct introduction into an aquifer. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

## Directions for Use

### Target Broadleaf Weeds, Application Rate, and Use Restrictions:

Target Broadleaf Weeds	Application Rate (pint/acre)	Use Restrictions
clover, white dandelion groundsel, common hawksbeard, smooth pineappleweed sowthistle, annual (suppression) thistle, bull (suppression) thistle, Canada (suppression) vetch	post-harvest application: 1/3 maximum  spring application: 1/3 maximum	<ul style="list-style-type: none"> <li>Do not exceed a total of 2/3 pint per acre (10.67 fl oz per acre; 0.25 lb ai per acre) per year.</li> <li>Do not apply within 4 months of planting.</li> <li><b>Postharvest Application:</b> Apply between September 15 and November 15.</li> <li><b>Spring Application:</b> Apply prior to first bloom. Do not apply within 30 days of harvest.</li> </ul>

### Broadcast Application

For difficult to control weeds, best results will be achieved with a post-harvest application followed by a spring application. Apply the spray uniformly in 30 to 75 gallons of water per acre using ground equipment only.

**Post-Harvest Application:** After harvest, make a ground application between September 15 and November 15 at a maximum rate of 1/3 pint of Stinger (0.125 lb ai clopyralid) per acre. For control of Canada thistle, apply Stinger after the majority of basal leaves have emerged but prior to bud stage.

**Spring Application:** Prior to first bloom, make a ground application in the spring at a maximum rate of 1/3 pint of Stinger per acre. Do not apply within 30 days of harvest.

**Spot Application:** Use a rate per acre equivalent to a broadcast application and follow the same use restrictions. Refer to the Hand-Held Sprayers section on the product label for Stinger for spot application instructions. Apply the spray uniformly.

### Specific Use Restrictions:

- **Chemigation:** Do not apply Stinger through any type of irrigation system.
- Do not apply Stinger by aircraft.
- Do not apply Stinger within 6 to 8 hours of expected rainfall or irrigation.
- Do not compost vegetation treated with Stinger.
- Do not use surfactants when applying Stinger to strawberry.
- Do not tank mix Stinger with other herbicides registered for use on strawberry.
- Refer to the product label for Stinger for crop rotation intervals.
- Do not apply Stinger to strawberry if you are unwilling to accept minor leaf injury. Minor leaf cupping may occur.

## **Special Conditions and Risks of Use**

**Dow AgroSciences Intends That This FIFRA Section 24(c) Special Local Need (SLN) Labeling be Used Only by Members of the Oregon Strawberry Commission.**

**USE OF Stinger<sup>®</sup> herbicide (THE "PRODUCT") ON STRAWBERRY (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.**

**Dow AgroSciences and the Oregon Strawberry Commission make the Product available for use in the manner described in this SLN Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this SLN Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.**

**By purchasing the Product for use, or using the Product, in the manner described in this SLN Labeling, you acknowledge and accept that:**

- (1) You assume all risk of Crop injury, Crop yield reduction and Crop loss;**
- (2) Dow AgroSciences and the Oregon Strawberry Commission do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this SLN Labeling;**
- (3) Dow AgroSciences and the Oregon Strawberry Commission do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;**
- (4) Dow AgroSciences and the Oregon Strawberry Commission disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;**
- (5) These Special Conditions and Risks of Use supersede any contrary representations or recommendations by Dow AgroSciences and the Oregon Strawberry Commission, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.**

**If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Dow AgroSciences and the Oregon Strawberry Commission and not specified by the U.S. EPA or the State of Oregon.**

24(c) Registrant:  
Dow AgroSciences LLC  
9330 Zionsville Road  
Indianapolis, IN 46268

<sup>®</sup><sup>TM</sup>Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow  
R043-082  
Accepted: 02/10/15  
Replaces R043-045