Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA



Weed Control Prior to the Emergence or Transplanting of Strawberries

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

Refer to the label booklet for this product for general information on product use, mixing, application equipment and techniques, labeled uses, and weed rate tables.

This product may be used to control annual and perennial weeds listed in the product label prior to emergence of strawberries. Applications should be made at least 3 days before transplanting or planting.

Precautions and Restrictions:

When applying this product prior to transplanting crops into plastic mulch, care must be taken to remove spray residues of this product from the plastic prior to transplanting; otherwise crop injury could occur. Removal of residues may be accomplished by application of 0.5 inch of water either by rainfall or sprinkler irrigation. Applications made at emergence will result in injury or death of emerged seedlings.

Up 48 fl oz/acre of this product may be applied.

[®]Trademark of Dow AgroSciences LLC R353-001 EPA accepted 04/23/07 Initial printing