

**RESTRICTED USE PESTICIDE****Toxic to fish and aquatic organisms**

For retail sale to and use only by certified applicators or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

(Not for use in New York or California)

SUPPLEMENTAL LABELING
FANFARE™ 2EC
INSECTICIDE/MITICIDE

EPA Reg. No. 66222-99

Contains bifenthrin, the active ingredient used in Capture 2EC.

Fanfare 2EC is not manufactured or distributed by FMC Corporation.

ACTIVE INGREDIENT:**% BY WT.**

Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl)methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-2,2-dimethyl-cyclopropanecarboxylate* 25.1%

OTHER INGREDIENTS:** 74.9%

TOTAL 100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.

**Contains petroleum distillates.

This product contains 2 pounds active ingredient per gallon.

Manufactured for:

Makhteshim Agan of North America, Inc.

4515 Falls of Neuse Rd., Suite 300

Raleigh, NC 27609

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product inconsistent with its labeling.
- This supplemental labeling must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- These new uses can be found on the EPA stamped label for Fanfare 2EC (66222-99) dated February 20, 2008.
- This supplemental label is not yet approved for use in the states of California and New York.

PREHARVEST INTERVAL

The required days to wait between the last application and harvest are given in () after each crop name.

CROPS**CHRISTMAS TREES (For Use Only in Washington and Oregon)**

PEST	DOSAGE		COMMENTS
	LB/AI/A	FL. OZ/A	
Root Weevil, Spruce Spider Mite	0.06-0.1	3.9-6.4	Ground application: Apply in water in a minimum of 20 gallons per acre. Air application: Apply in water in a minimum of 5 gallons per acre. Fanfare is generally not phytotoxic to Christmas trees. However, applications should be made to a small representative group of plants to ensure that a particular variety grown under current conditions is not unusually sensitive to Fanfare 2EC. Maintain a minimum of 21 days between applications.
<ul style="list-style-type: none">• Do not apply more than 0.1 lb. active ingredient (6.4 ounces formulated) per acre per treatment.• Do not make more than 3 applications in a crop year.• Do not make applications through irrigation systems.			

CILANTRO, CORIANDER (3)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Beet Armyworm, Cabbage Looper, Cutworm, Flea Beetle, Grasshoppers, Leafminer, Saltmarsh Caterpillar, Spotted Cucumber Beetle, Thrips, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain thorough coverage.
Two Spotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season.Do not make applications less than 7 days apart.			

CONIFER SEED ORCHARDS

(For Use Only in Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Oklahoma, South Carolina, Tennessee, Texas, Virginia)

PEST	DOSAGE		REMARKS
	LB/AI/A	FL. OZ/A	
Cone Worms, Seed Bugs, Seed Worms	0.1-0.2	6.4-12.8	Ground application: Apply in water in a minimum of 100-500 gallons per acre. Air application: Apply in water in a minimum of 10 gallons per acre or 0.5 gallon refined vegetable oil per acre. Apply in sufficient water to obtain thorough coverage. Begin applications 7 days after peak pollen flight and continue on 30 day intervals up to a maximum of 0.6 lb. active per acre per season.
<ul style="list-style-type: none"> Do not make more than six applications per season or apply more than 0.6 lb. active ingredient (38.4 ounces formulated) per acre per season. 			

DRIED BEANS AND PEAS (14)

Dried cultivars of:

Bean (*Lupinus*): Grain Lupin, Sweet Lupin, White Lupin, White Sweet Lupin

Bean (*Phaseolus*): Field bean, Kidney Bean, Lima Bean (dry), Navy Bean, Pinto Bean, Tepary Bean

Bean (*Vigna*): Adzuki Bean, Blackeyed Pea, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean

Broad Bean (dry); Chickpea; Guar; Lablab Bean; Lentil; Pea (*Pisum*), Field pea; Pigeon Pea

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Banks Grass Mite, Twospotted Spider Mite, Carmine Mite, Lygus spp.	0.08-0.10	5.12-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. Thorough coverage is essential to achieve control.
Aster Leafhopper, Flea Beetle, Grasshoppers, Leafhoppers	0.025-0.10	1.6-6.4	
Aphids, Beet Armyworm, Fall Armyworm, Southern Armyworm, Yellowstriped Armyworm, Bean Leaf Beetle, Cucumber Beetles, Japanese Beetle, Adult Sap Beetle, Plant Bug, Stink Bugs, Tarnished Plant Bug, Alfalfa Caterpillar, Cloverworm, European Corn Bore*, Cutworms, Western Bean Cutworm, Corn Earworm, Loopers, Corn Rootworm Adults, Thrips, Webworms, Pea Weevil, Pea Leaf Weevil, Whitefly, Imported Cabbageworm, Saltmarsh Caterpillar, Tobacco Budworm, Leafminer	0.33-0.10	2.1-6.4	
<ul style="list-style-type: none">Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) to beans per acre per season.Do not make applications less than 7 days apart.			

LEAFY BRASSICAS (7)

Broccoli Raab, Bok Choy, Kale, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Crickets, Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles, Grasshoppers, Ground Beetles, Imported Cabbageworm, Japanese Beetle (adult), Leafhoppers, Loopers, Saltmarsh Caterpillar, Stink Bugs, Thrips, Tobacco Budworm, Whitefly, Wireworm (adults)	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Emulsified oil may be substituted for water. See section entitled MIXING INSTRUCTIONS for details on the amount of oil to use in the spray tank in lieu of water. Thorough coverage is essential to achieve control.
Banks Grass Mite, Carmine Mite, Lygus spp., Pacific Spider Mite, Twospotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">Do not apply more than 0.4 lb. active ingredient (25.6 ounces formulated) per acre per season.Repeat applications if needed to maintain control, but do not make applications less than 7 days apart.			

OKRA (7)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Aphids, Armyworms, Corn Earworm, Cucumber Beetles, Cutworms, European Corn Borer, Flea Beetles, Japanese Beetle (Adult), Leafminers, Loopers, Stink Bugs, Thrips, Whitefly	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons of finished spray per acre. Air application: Apply in water in a minimum of 2 gallons per acre. Apply in sufficient water to obtain uniform coverage as needed.
Broad Mite, Carmine Mite, Lygus spp., Two Spotted Spider Mite	0.08-0.10	5.12-6.4	
<ul style="list-style-type: none">• To maintain a proper spray interval, do not make applications less than 7 days apart.• Do not apply more than 0.20 lb. active ingredient (12.8 ounces formulated) per acre per season.			

TOBACCO

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Armyworm spp., Cutworm spp., Mole Crickets, Stalkborers, Tobacco Flea Beetle (larvae), White Grubs, Wireworms	0.625-0.10	4.0-6.4	Pre-transplant soil applications: Apply 0.0625-0.1 active ingredient per acre in a minimum of 10 gallons per acre to control soil pests. Use of suitable equipment to incorporate into top 4" of the soil is required to control below ground pests. At-transplant water treatment application: Apply 0.0625-0.1 lb. active ingredient per acre in a water treatment application volume of the 10-200 gallons per acre. May be tank mixed with Command®, Spartan®, and other herbicides approved for tobacco use. Foliar applications: Apply 0.04-0.10 lb. active ingredient per acre foliar application up to and including layby in a minimum of 10 gallons per acre. May be tank mixed with Command, Spartan, and other herbicides approved for tobacco use.
Aphid spp., Armyworm spp., Chinch Bugs, Cutworm spp., Flea Beetle (adults), Grasshoppers, Green Bugs, Japanese Beetles, Stink Bugs, Tarnished Plant Bugs, Thrips, Whiteflies	0.04-0.10	2.56-6.4	
Lygus spp., Spider mites	0.1	6.4	
<ul style="list-style-type: none">For foliar applications, do not make more than 2 applications per season.For all applications, do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.Do not apply later than layby.			

TUBEROUS AND CORM VEGETABLES (21)

Arracacha; Arrowroot; Chinese Artichoke; Jerusalem Artichoke; Edible Canna; Cassava (bitter and sweet); Chayote (root); Chufa; Dasheen (taro); Ginger; Leren; Potato; Sweet Potato; Tanier; Tumeric; Yam Bean; True Yam

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Corn Wireworm, Tobacco Wireworm	0.30 (at plant)	19.2 (at-plant)	In-furrow planting time treatment: Fanfare 2EC may be applied as an in-furrow planting time treatment for the control of wireworms, rootworms, and white grubs. Apply Fanfare 2EC at the rate of 0.3 pounds active ingredient per acre as an in-furrow spray or T-band spray at planting time. Lay-by treatment: Fanfare 2EC may be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Fanfare 2EC to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Fanfare 2EC as a banded spray over the row at a rate of 0.05-0.15 pounds active ingredient per acre (3.2 to 9.6 ounces formulated) in 10 gallons per acre of spray. Foliar spray: Fanfare 2EC may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles, and May/June beetles (white grubs). Apply Fanfare 2EC at the rate of 0.1 lb. active ingredient per acre (6.4 ounces formulated) in 10 gallons of spray by ground and 3 gallons of spray by air.
Southern Potato Wireworm, Japanese Beetle Grubs, June Beetle	0.05-0.15 (lay-by)	3.2-9.6 (lay-by)	
Sweetpotato Flea Beetle, Cucumber Beetle, Sweetpotato Weevil, Banded Cucumber Beetle, Black Flea Beetle, Whitefringed Beetle, White Grub, Sugarcane Beetle, Rootworms	0.033-0.10 (foliar)	2.1-6.4 (foliar)	
<ul style="list-style-type: none">For foliar applications, do not make more than 2 foliar applications per season and do not make application less than 21 days apart.Do not apply more than 0.5 lb. active ingredient (32 ounces formulated) per acre per season, including soil applications.			

LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Fanfare 2EC (66222-99)(EPA app 2-20-08)-supplemental