

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Entrust[®]

EPA Reg. No. 62719-282

Control of Lepidopteran Larvae and Suppression of Flea Beetles, Dipteran Leafminers and Thrips in Spices¹ (Except Black Pepper) (Subgroup 19B)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Entrust[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Entrust according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Entrust.

Directions for Use

¹Spices, including: allspice, anise (seed), annatto (seed), black caraway, caper (buds), caraway, cardamom, cassia (buds), celery (seed), cinnamon, clove (buds), common fennel, coriander (seed), culantro (seed), cumin, dill (seed), Florence fennel (seed), fenugreek, grains of paradise, juniper (berry), lovage (seed), mace, mustard (seed), nutmeg, poppy (seed), saffron, star anise, vanilla, white pepper

Refer to product label for Entrust for General Use Precautions, Mixing and Application instructions.

Pests and Application Rates:

Pests	Entrust (oz/acre)	Restrictions
lepidopteran larvae	1.25 - 2	<ul style="list-style-type: none">• Preharvest Interval: Do not apply within 14 days of harvest.• Re-treatment Interval: 10 days• Do not apply more than a total of 9 oz of Entrust (0.45 lb ai of spinosad) per acre per crop or make more than 5 applications per calendar year.
flea beetles dipteran leafminers ¹ thrips ¹	2 - 3	

¹Suppression of leafminers and thrips may be improved with the addition of an adjuvant to the spray mixture. See Use of Adjuvants under Mixing.

Specific Use Directions:

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated to control target pests. Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: For resistance management purposes, do not apply Entrust more than 3 times in any 30-day period or apply sprays less than 10 days apart. Whenever Entrust is applied up to 3 times in succession, this should be followed by no use of Entrust for a 30-day period or rotation to another insecticide class.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R184-022

Accepted: 02/27/08

Initial printing.