

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Disease Control on Black Sapote, Canistel, Mamey Sapote, Mango, Papaya, Sapodilla, and Star Apple

Refer to product label for General Use Directions, Mixing and Application instructions.

Disease	Rally 40WSP (oz/acre)	Application Timing	Restrictions
powdery mildew (<i>Oidium caricae</i>)	10	Scout crops on a regular basis and treat when disease first appears or when conditions favor disease development. Apply uniformly in a spray volume that provides thorough coverage of the fruit and foliage. Control may be reduced at low spray volumes or if spray coverage is not adequate. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.	<ul style="list-style-type: none"> • Do not apply more than 10 oz (0.63 lb) of Rally 40WSP (0.25 lb ai) per acre per application or 80 oz (5 lb) of Rally 40WSP (2 lb ai) per acre per season. • Do not make more than 8 applications per season. • Pre-harvest Interval: None. • Re-treatment Interval: 14 days • Rotational Crop Restriction: Do not plant any crop other than those listed on the main label for 12 months following the last application of Rally 40WSP.

Resistance Management: Rally 40WSP belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Rally 40WSP should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

®Trademark of Dow AgroSciences LLC

R182-001

EPA accepted 03/20/08

Initial printing