



FANFARE® 2EC
EPA REG. NO. 66222-99

**FIFRA 2(ee) RECOMMENDATION
FOR CONTROL OF TOBACCO BUDWORM IN TOBACCO**
For use in North Carolina, South Carolina, and Tennessee Only

DIRECTIONS FOR USE

This FIFRA Section 2(ee) Recommendation recommends using Fanfare 2EC on tobacco for control and/or suppression of listed pests. The user must refer to the federally approved Fanfare 2EC label and read and follow all directions for use, restrictions, and precautions. This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. It is a violation of federal law to use this product in a manner inconsistent with its labeling. Users should have this recommendation in their possession at the time of use.

TOBACCO

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Tobacco Budworm ¹	0.04-0.10	2.56-6.4	Foliar applications: Apply 0.04-0.10 lb. active ingredient per acre foliar application up to and including layby in a minimum of 10 gallons per acre. May be tank mixed with Command, Spartan, and other herbicides approved for tobacco use.
<ul style="list-style-type: none">• For foliar applications, do not make more than 2 applications per season.• For all applications, do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.• Do not apply later than layby.			

¹ Refer to **RESISTANCE MANAGEMENT** statement on the label attached to the container for specific information regarding resistance management with this product and this pest.