

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

SpinTor[®] 2SC
EPA Reg. No. 62719-294

EPA 24(c) Special Local Need Registration SLN SC-080005
For Distribution and Use Only Within the State of South Carolina

Suppression of Thrips in Ginkgo

Special Conditions and Risks of Use

USE OF SpinTor[®] 2SC Naturalyte[®] insect control (THE "PRODUCT") ON GINKGO (THE "CROP") MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS. READ AND UNDERSTAND THESE SPECIAL CONDITIONS AND RISKS OF USE BEFORE USING THE PRODUCT ON THE CROP.

Dow AgroSciences makes the Product available for use in the manner described in this Supplemental Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury or loss. The decision to use this Product in the manner described in this Supplemental Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls, and (iv) any other relevant factors.

By purchasing the Product for use, or using the Product, in the manner described in this Supplemental Labeling, you acknowledge and accept that:

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental Labeling;
- (3) Dow AgroSciences does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaims all warranties, express or implied, including any implied warranties of fitness for a particular purpose or merchantability;
- (4) Dow AgroSciences disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Special Conditions and Risks of Use supersede any contrary representations or recommendations by Dow AgroSciences, its agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Special Conditions and Risks of Use are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container. These Special Conditions and Risks of Use are required by Dow AgroSciences and not specified by the U.S. EPA or the State of South Carolina.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for SpinTor® 2SC Naturalyte® insect control before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of SpinTor 2SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for SpinTor 2SC.

Directions for Use

Pest	SpinTor 2SC fl oz/acre
thrips	5 – 10 (0.078 – 0.156 lb ai/acre)

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size. Base your application timing upon local thresholds. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply as a foliar spray at the rate indicated for thrips in enough spray volume to get thorough coverage. Use application equipment suitable for insecticide spraying for this crop. Use a higher rate in the rate range for heavier infestations. Thorough coverage is required for optimum efficacy. Prolonged or heavy infestations may require repeat applications. Suppression of thrips may be improved by addition of an adjuvant approved for growing crops to the spray mixture.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 10 days of harvest.
- Do not apply more than a total of 29 fl oz of SpinTor 2SC (0.45 lb ai spinosad) per acre per growing season.
- **Chemigation:** Do not apply this product to ginkgo through any type of irrigation system.
- Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least 2 applications.
- Do not harvest plants for food or feed for at least one year after treatment.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R064-010

Accepted: 05/06/08

Initial printing.