

**SUPPLEMENTAL LABELING  
FOR DISTRIBUTION AND USE ONLY WITHIN KENTUCKY**



# **Terramaster® 4EC**

**FUNGICIDE**

**FOR USE ON GREENHOUSE TOMATOES**

**EPA REG. NO. 400-422      EPA SLN NO. KY-080004**

Active Ingredient: (% by weight)

Etridiazole [5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole]\* ..... 44.3%

Inert Ingredients:\*\* ..... 55.7%

Total: ..... 100.0%

\*Contains 4 lbs. of 5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole per U.S. Gallon at 20°C.

\*\*Contains Petroleum Distillates

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)**

**DIRECTIONS FOR USE**

It is a violation of State and Federal law to use this product in a manner inconsistent with its labeling.

**GREENHOUSE TOMATOES**

**For control of Pythium and Phytophthora Root Rot in  
greenhouse tomatoes by drip irrigation**

**Drip Irrigation**

Apply TERRAMASTER 4EC in a 0.01% solution (6.5 fluid oz. in 500 gallons of water) no sooner than 3 weeks after transplanting, using a volume of 6 - 8 fl. oz. per plant.

Reapply as needed, but no sooner than 3 weeks after a previous application.

Do not exceed 4 applications of TERRAMASTER 4EC per crop.

Do not apply solutions more concentrated than 0.01% without first verifying potential phytotoxicity on the variety of tomato being grown.

Do not apply TERRAMASTER 4EC sooner than 3 days before harvest.

Uniform distribution of TERRAMASTER 4EC through the growing medium is essential for best results. Localized over-dosing may cause crop damage and under-dosing may cause ineffective disease control.

TERRAMASTER 4EC should be introduced into the growing medium only after the medium has been thoroughly irrigated to ensure adequate product movement through the growing medium. Continue irrigation until the entire desired quantity of TERRAMASTER 4EC has been introduced.

Do not mix TERRAMASTER 4EC with other pesticides or fertilizers unless compatibility tests have shown complete compatibility.

Apply TERRAMASTER 4EC to a small crop area to ensure crop safety before applying to broad crop areas.

**THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER**

**Follow all applicable directions, restrictions, and precautions on the EPA registered label.**

**This label must be in the possession of the user at the time of the pesticide application.**

**001**

Terramaster® is a Registered Trademark of Chemtura Corporation

24c (SLN) REGISTRANT:

Chemtura Corporation

199 Benson Road

Middlebury, CT 06749

EPA SLN NO. KY-080004