

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Starane<sup>®</sup> Ultra

EPA Reg. No. 62719-577

### Weed Control in Millet Grown for Grain, Forage or Hay

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane<sup>®</sup> Ultra herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Starane Ultra according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Starane Ultra.

#### Directions for Use

**Starane Ultra may be applied for broadleaf weed control in millet grown for grain, forage or hay.**

Apply Starane Ultra as a broadcast postemergence treatment using ground equipment or by air. A second application may be made a minimum of 14 days after the first. Starane Ultra may be applied in tank mix combination at labeled rates with other herbicides registered for these uses. All applicable use directions, precautions and limitations on the labels of the tank mix products must be followed. When tank mixing, the most restrictive limitations on each label must apply.

**Application Timing:** Apply to millet in the spring when weeds are actively growing and before weeds are 8 inches tall. Only weeds emerged at the time of treatment will be controlled. New plantings of millet may be treated from the 2 true leaf stage of growth prior to early boot stage. Do not apply during boot, flowering, or seed development stage of growth if grass crop is to be harvested for seed.

**Broadcast Application Rates:** (Numbers in parentheses (-) refer to footnotes following table.)

Weed Size or Species (1)	Starane Ultra (pt/acre)
Susceptible broadleaf weed seedlings less than 4 inches tall (2)	0.3
Susceptible broadleaf weed seedlings less than 8 inches tall or vining	0.4

1. Refer to the Weeds Controlled or Suppressed section in the label booklet for Starane Ultra for a complete listing of weeds controlled or suppressed.
2. The 0.3 pint/acre rate will generally provide satisfactory control of kochia seedlings less than 4 inches tall (including ALS resistant biotypes). However, when conditions for control are less favorable, such as under drought or cool temperatures, the 0.4 pint/acre rate will provide more consistent control of kochia seedlings 1 to 4 inches tall. Control of small kochia with reduced rates will be more consistent if kochia is at least 1 inch tall. The 0.4 pint/acre rate should be used for optimal control of dicamba tolerant kochia populations (see Management of Kochia Biotypes in the General Information section of the label booklet for Starane Ultra.

**Restrictions:**

- Do not apply more than 0.7 pint per acre of Starane Ultra or make more than 2 applications per growing season.
- **Grazing restrictions:** There are no grazing restrictions for lactating or non-lactating dairy animals.
- **Preharvest Interval:** Do not harvest millet hay within 14 days of application or millet grain and straw within 40 days of application.
- **Slaughter restrictions:** Meat animals must be withdrawn from treated forage at least 2 days before slaughter.

®Trademark of Dow AgroSciences LLC

R357-006  
EPA accepted 06/10/08  
Initial printing.