


**RESTRICTED USE PESTICIDE.**

Due to acute toxicity,  
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



**FIFRA Section 24(c) Supplemental Label  
FIRESTORM®**

**FOR DISTRIBUTION AND USE  
ONLY WITHIN THE STATE OF TEXAS**

**EPA Reg. No. 82557-1-400**

**EPA SLN No. TX-080020**

**FOR HARVEST AID APPLICATIONS ON COTTON IN TEXAS  
(EXCEPT IN THE COUNTIES OF STARR, HIDALGO, WILLACY, AND CAMERON)**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered product label.
- This label must be in the possession of the user at the time of application.

Failure to follow the directions for use and precautions on this label may result in poor pest control, crop injury, or illegal residues.

**DIRECTIONS FOR USE: Harvest Aid Applications on Cotton in Texas (except in the counties of Starr, Hidalgo, Willacy, and Cameron)**

Up to three multiple broadcast harvest aid applications of FIRESTORM® may be used. **Do not exceed a total of 32 fluid ounces (fl. oz.) (0.75 lb. active ingredient) per acre per crop season.** If multiple applications are made, allow 7 days between applications. Do not apply to cotton within 3 days of harvest. Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Crop	FIRESTORM® Rate per Acre	Precautions, Restrictions and Comments
Tank mixes with phosphate or chlorate defoliant	5.4 - 10.7 fl. oz. (0.13 to 0.25 lb a.i.) alone <u>or</u> tank mixed with 1½ - 2 gal. chlorate <u>or</u> 1½ - 2 pints phosphate	<ul style="list-style-type: none"> <li>• Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.</li> <li>• In desert cotton areas or on rank vigorous cotton, use the 10.7 fl oz. (0.25 lb a.i.) rate of FIRESTORM.</li> <li>• Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Follow the Harvest Aid Use Precautions listed on the Federal label.</li> </ul>
Tank mixes for boll opening and defoliation of Southern cotton	2.7 to 3.3 fl. oz. (0.06 to 0.08 lb a.i.)	<ul style="list-style-type: none"> <li>• To aid in defoliation and opening of mature bolls, FIRESTORM may be tank mixed with the following products: DEF® defoliant, Dropp® defoliant, Prep™ PGR, Folex® defoliant, Accelerate® defoliant</li> <li>• Apply when 60% or more of the bolls are open and remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Refer to tank mix product label for rate, directions, limitations and precautions.</li> </ul>
Harvest aid for defoliation and boll opening of stripper or spindle-harvested cotton	2.1 to 7.5 fl. oz. (0.05 to 0.18 lb a.i.)	<ul style="list-style-type: none"> <li>• Apply when <b>75%</b> or more of the bolls are open and remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• FIRESTORM may be applied alone or tank mixed with the cotton harvest aid products listed above.</li> <li>• To avoid leaf sticking, apply FIRESTORM as a desiccant approximately 3 to 7 days after defoliation or a conditioning application and 7 to 14 days before harvest.</li> <li>• Cooler temperatures may cause longer waiting period between application of FIRESTORM as a desiccant and a subsequent defoliation/conditioning treatment.</li> <li>• Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Late season harvest aid desiccation of stripper cotton or spindle-harvested cotton	11.2 to 21 fl. oz. (0.26 to 0.49 lb a.i.)	<ul style="list-style-type: none"> <li>• Use to desiccate regrowth occurring after defoliation or desiccation.</li> <li>• Apply when <b>85%</b> of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• To avoid leaf sticking, apply FIRESTORM as a desiccant approximately 3 to 7 days after defoliation or a conditioning application and 7 to 14 days before harvest.</li> <li>• Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Desiccation of regrowth	11.2 to 21 fl. oz. (0.26 to 0.49 lb a.i.)	<ul style="list-style-type: none"> <li>• Use to desiccate regrowth occurring after defoliation or desiccation.</li> <li>• Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary.</li> <li>• Control is dependent upon growing condition and desiccation of small new growth may not always be complete.</li> <li>• Use higher rate if regrowth is excessive.</li> </ul>

Refer to the FIRESTORM federal label for additional cotton recommended uses and nonionic surfactant recommendations.

**Additional Precautions, Restrictions and Comments:**

- Because of extremes in environmental and plant conditions, it is advisable to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliants, observe livestock precautions listed on the phosphate label.
- Do not apply to cotton within 3 days of harvest. Do not harvest within 7 days of application when used with phosphate or chlorate defoliants.

**Do not apply this product through any type of irrigation system.**

**This bulletin contains new or supplemental instructions for use of the product which may not appear on the package label.**

**24(c) REGISTRANT:**

Sinon USA Inc.  
1080 Carol Lane, Suite 264  
Lafayette, CA 94549

Distributed by:  
Chemtura Corporation  
199 Benson Road  
Middlebury, CT 06749

FIRESTORM® is a registered trademark of Chemtura Corporation  
ACCELERATE® is a registered trademark of Cerexagri, Inc.  
DEF®, DROPP®, PREPTM are trademarks of Bayer CropScience  
FOLEX® is a registered trademark of AMVAC Chemical Corporation