

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA 24(c) Special Local Need Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban[®] Advanced

EPA Reg. No. 62719-591

**EPA 24(c) Special Local Need Registration SLN CA-080010
For Distribution and Use Only in the State of California**

Control of Argentine Ants Infesting Grapes

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Apply 1.5 to 2 pints of Lorsban Advanced per acre in a minimum of 2.5 gallons of water per acre using a power-operated ground or hand-held sprayer. Maintain constant agitation of the spray tank. To obtain thorough coverage, spray the base of each vine, the vine stakes, and the surrounding soil to about 1 foot from the base of the vine. If a hand-held sprayer is used, 2 to 5 fl oz per vine is required. A non-petroleum-based spreader/sticker spray adjuvant may be used if previous experience indicates that it is safe under the intended conditions of use. Follow use directions for the spreader/sticker product.

Specific Use Precautions:

- Do not allow spray to contact fruit or foliage.
- Weed growth in the vine rows should be controlled to prevent ants from trailing to the vine cordons via weeds and to allow better spray coverage.
- Consult your extension service specialist or certified pest control advisor for use information specific to your area.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 76 days before harvest.
- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than 3 applications per year.
- Maximum single application rate is 0.94 lb ai chlorpyrifos per acre

- For residue purposes, growers may treat grapes using Lorsban Advanced under the requirements of this SLN or Lorsban Advanced under the requirements listed in SLN CA-080009. Growers may not treat the same grape crop under the requirements of both SLNs, nor may they treat the same grape crop under the requirements of any SLN for Lorsban-4E.

Expiration Date: December 31, 2019

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R368-105

Approved: 10/13/14

Replaces R368-050