

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®] Advanced

EPA Reg. No. 62719-591

Insect Control for Onion (Dry Bulb)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

At-Plant Soil Drench Application

For direct seeded onions to control onion maggot, apply 32 fl oz of Lorsban Advanced per acre in a water-based spray as a 2- to 4-inch wide band over the row at planting time in a minimum of 40 gpa. Equivalent rates of insecticide spray required per 1000 feet of row for listed row spacings are given in the accompanying table. Shallow incorporation is necessary. Place behind the planter shoe and in front of the presswheel. Phytotoxicity may occur if Lorsban Advanced is sprayed directly onto onion seeds. Do not mix Lorsban Advanced with other pesticide products. **Note:** The user should exercise reasonable judgment and caution with this product. Until familiar with results under user planting and growing conditions, limit application of this product to a small area to determine plant tolerance and extent of injury if such occurs prior to initiating large scale applications.

Lorsban Advanced (32 fl oz/acre)	Row Spacing			
	6-inch	10-inch	12-inch	18-inch
fl oz/1000 ft of row	0.37	0.61	0.74	1.1

Specific Use Restrictions:

- Do not make more than one application per year.
- Maximum single application rate is 0.032 lb ai chlorpyrifos per 1000 feet of row.
- Do not aerially apply this product in Mississippi.

Postplant Soil Drench Application

Apply as an early season directed spray to the base of onion seedlings or transplants during peak egg laying. Use a minimum of 100 gpa for thorough wetting.

Target Pests	Lorsban Advanced (quart/acre)
onion maggot seedcorn maggot	1

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 60 days before harvest.
- Do not make more than two applications (at plant plus postplant) per year.
- Maximum single application rate is 0.94 lb ai chlorpyrifos per acre.
- Do not aerially apply this product in Mississippi.

®Trademark of Dow AgroSciences LLC

R368-007

EPA accepted: 06/06/08

Initial printing.