

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®] Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN NV-080004
For Distribution and Use Only in the State of Nevada

Control of Billbugs, Cutworms and Aphids Infesting Perennial Grass Seed Crops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Billbugs and Cutworms: Apply 1 quart of Lorsban Advanced per acre in 20 gallons of water per acre as a broadcast spray using power-operated ground spray equipment. For improved results, thatch should be moist at time of application and the insecticide should be washed into the thatch immediately after application. If sprinkler irrigation is available, apply 1/4 to 1/2 inch of water after application to help incorporate the material, or apply to early morning dew-covered fields, or during or immediately following a light rain. For **billbug control**, apply in early April when adult billbugs are active. For **cutworm control**, apply in either spring or fall when larvae or larval damage first appears. Fall applications should follow sufficient irrigation and/or rainfall. Heavy carbon residue from field burning will prevent Lorsban Advanced penetrating into the plant crown where the insects feed. Re-treat as necessary to maintain control.

Aphids: Apply 1 to 2 pints of Lorsban Advanced per acre in 20 gallons of water per acre as a broadcast spray using power-operated ground spray equipment. Treat before heading as colonies begin to form.

Consult your cooperative agricultural extension service for additional information concerning billbug, cutworm, and aphid control practices in your area.

Specific Use Precautions and Restrictions:

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than three applications of Lorsban Advanced per year.
- Maximum single application rate is 0.94 lb ai chlorpyrifos per acre.
- Apply only to perennial grass seed crops.
- Do not allow meat or dairy animals to graze in treated areas.
- Do not feed treated grass cuttings or seed screenings to livestock or use hay for livestock bedding.

- This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your cooperative agricultural extension service.

®Trademark of Dow AgroSciences LLC

R368-033

Approved: 11/19/08

Initial printing.