

**FIFRA SECTION 24(c) SUPPLEMENTAL LABELING  
FOR DISTRIBUTION AND USE ONLY WITHIN CALIFORNIA**

**RESTRICTED USE PESTICIDE**

**Due to eye and skin effects.**

For retail sale to and use by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



**Comite®**

**AGRICULTURAL MITICIDE**

**EPA REG. NO. 400-104**

**COMPOSITION**

Active Ingredients: (% by weight)

Propargite (Sulfurous acid, 2-[4-(1,1-dimethyl-ethyl) phenoxy] cyclohexyl-2- propynyl ester)\* .....73.6%

Other Ingredients .....26.4%

Total: ..... 100.0%

\*Contains 6.55 lbs. active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN  
DANGER • PELIGRO**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

**NON-BEARING ALMONDS AND WALNUTS INTERPLANTED WITH DRY BEANS AND COTTON:**

For the control of two-spotted spider mites.

**GROUND APPLICATION:**

**FOR DRY BEANS:** Use 2 to 3 pints of COMITE per acre in a minimum of 25 gallons of water per acre. Do not exceed 2 applications per year. Do not apply more than 4.5 pints of COMITE per acre per season, equivalent to 3.7 lbs propargite a.i. per acre per season. Observe an REI of 21 days. Allow a minimum of 21 days between any 2 applications. The earliest harvest days is 14 days after the last application.

**FOR COTTON:** Use 2 pints of COMITE per acre in a minimum of 25 gallons of water per acre. Do not exceed 2 applications per year. Do not apply more than 4.0 pints of COMITE per acre per season, equivalent to 3.3 lbs propargite a.i. per acre per season. Applications can be made up to boll opening. Observe an REI of 7 days. Allow a minimum of 21 days between any 2 applications.

**AERIAL APPLICATION:**

**FOR DRY BEANS:** Use 2 to 3 pints of COMITE per acre in a minimum of 10 gallons of water per acre. Do not exceed 2 applications per year. Do not apply more than 4.5 pints of COMITE per acre per season, equivalent to 3.7 lbs propargite a.i. per acre per season. Observe an REI of 21 days. Allow a minimum of 21 days between any 2 applications. The earliest harvest days is 14 days after the last application.

**FOR COTTON:** Use 2 pints of COMITE per acre in a minimum of 10 gallons of water per acre. Do not exceed 2 applications per year. Do not apply more than 4.0 pints of COMITE per acre per season, equivalent to 3.3 lbs propargite a.i. per acre per season. Applications can be made up to boll opening. Observe an REI of 7 days. Allow a minimum of 21 days between any 2 applications.

COMITE is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces is necessary for effective control.

**Follow all applicable directions, restrictions, and precautions on the EPA registered label.**

**This label must be in the possession of the user at the time of the pesticide application.**

**005**