

FIFRA Section 24(c) Special Local Need

(SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally® 40WSP

EPA Reg. No. 62719-410

EPA SLN No. OR-090006

For Distribution and Use Only Within the State of Oregon

This label valid until December 31, 2019 or until otherwise amended, withdrawn, canceled, or suspended.

Control of Rusts (*Puccinia* spp.) and Powdery Mildew (*Erysiphe graminis*) in Perennial Grasses Grown for Seed Including Perennial Ryegrass, Fescue, Orchardgrass and Kentucky Bluegrass

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN labeling and the federal label for this product must be in the possession of the user at the time of application.
- Read this SLN label and the label affixed to the container for Rally® 40WSP fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Rally 40WSP according to this SLN label is subject to all use precautions and limitations imposed by the label affixed to the container for Rally 40WSP.

Active Ingredient

myclobutanil: a-butyl-a-(4-chlorophenyl)-1H-
1,2,4, triazole-1-propanenitrile40%

Other Ingredients60%

Total100%

Keep Out of Reach of Children

WARNING

Environmental Hazards

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

Directions for Use

Rally® 40WSP fungicide is a systemic curative and protectant fungicide. For best results, apply on a regularly scheduled protectant spray schedule, starting at the first sign of disease development and continuing until the threat of further disease development is past. Apply 5 to 7.5 oz (0.125 to 0.187 lb active) of Rally 40WSP in a minimum of 20 gallons of water per acre for ground application or in a minimum of 10 gallons of water per acre for aerial application. Begin applications in early spring or when disease symptoms are first noticeable and continue applications on a 14- to 21-day application schedule. Use a higher rate or shorter spray interval on susceptible varieties or under heavy disease pressure.

Grass Species Grown for Seed		Target Pests	
fescues	<i>Festuca</i> spp.	rusts	<i>Puccinia</i> spp.
orchardgrass	<i>Dactylis</i> spp.		
perennial ryegrass	<i>Lolium</i> spp.		
Kentucky bluegrass	<i>Poa pratensis</i>	powdery mildew	<i>Erysiphe graminis</i>

Specific Use Restrictions:

- Do not harvest grass hay from treated fields within 1 year (365 days) after the last application of Rally 40WSP.
- Do not graze livestock in treated fields within 1 year (365 days) after the last application of Rally 40WSP.
- Do not apply more than 30 oz product (0.75 lb active) per acre per year.
- **Chemigation:** For use under this SLN label, do not apply this product through any type of irrigation system.
- Do not cut treated grass grown for seed for forage, hay, or silage.
- Do not use harvested seed for sprouting.
- No portion of the treated field, including seed, seed screenings, hay forage or stubble may be used for human or animal feed.

24(c) Registrant:
Dow AgroSciences, LLC
9330 Zionsville Road
Indianapolis, IN 46268

®Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

R182-065
Accepted: 03/03/15
Replaces: R182-032