

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban[®] Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN VA-090001
For Distribution and Use Only in the State of Virginia

Insect Control Infesting Tobacco

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban[®] Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Directions for Use

Refer to product label for Lorsban Advanced for Use Precautions and Restrictions, Mixing and Application instructions.

Apply as a preplant broadcast spray to reduce the feeding damage caused by listed pests. Apply 24 to 48 hours before bedding and transplanting using a spray volume of 10 gallons per acre (gpa) or more. Incorporate immediately after application to a depth of 2 to 4 inches using suitable incorporation equipment.

Before broadcast application of Lorsban Advanced onto existing beds, knock down beds to final shape for transplanting. Use of PTO-driven implements that will incorporate Lorsban Advanced to a depth of 4 inches is recommended.

Target Pests	Lorsban Advanced (quart/acre)
cutworms flea beetles mole crickets root maggots wireworms	2

To control the above listed pests and suppress populations of rootknot nematodes in all tobacco growing regions, use Lorsban Advanced in a tank mix with Nemacur 3 at the rate of 2 quarts of Lorsban Advanced

plus 4 quarts of Nematicur 3 nematicide per acre. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for Nematicur 3 used in combination with Lorsban Advanced.

Apply the specified rate(s) to the soil surface in a spray volume of 10 gpa or more 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment. Where the nematode species *Meloidogyne arenaria* or *M. javanica* are present or high populations of *M. incognita*, apply Telone® II soil fumigant at the recommended label rate.

Specific Use Precautions and Restrictions:

- **Worker Restricted Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.
- Do not make more than one application of Lorsban Advanced or other product containing chlorpyrifos per season.
- Maximum single application rate is 1.88 lb ai chlorpyrifos per acre.
- **Chemigation:** Do not apply through any type of irrigation equipment.

®Trademark of Dow AgroSciences LLC

R368-047

Approved: 04/01/09

Initial printing.