

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Delegate™ WG

EPA Reg. No. 62719-541

Foliar Insect Control in Hops, Dried Cones

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Delegate™ WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Delegate WG according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate WG.

Directions for Use

Refer to product label for Delegate WG for General Use Precautions, Mixing and Application instructions.

Pests and Application Rates:

Pests	Application Rate		Recommendations and Restrictions
	Active Ingredient (lb/acre)	Delegate WG (oz/acre)	
lepidopterous larvae including: armyworms cutworms leafrollers loopers thrips (suppression)	0.039 – 0.063	2.5 – 4	Use Restrictions: <ul style="list-style-type: none">• Do not apply more than a total of 19.5 oz of Delegate WG (0.305 lb ai of spinetoram) per acre per crop.• Maximum Number of Applications: Do not make more than 5 applications per crop.• Minimum Re-Treatment Interval: Do not make applications less than 4 days apart.• Preharvest Interval (PHI): Do not apply within 1 day of harvest.

Specific Use Directions:

Application Timing: For determining when to treat, scout with enough regularity to monitor the population size of each of the labeled pests. Treat when pests appear, targeting eggs at hatch or small larvae. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Application Rate: Apply Delegate WG as a foliar spray at the rate indicated to control target pests. Use a higher rate in the specified rate range for larger larvae or high infestations and/or larger plant volume. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: Do not make more than 2 consecutive applications of Group 5 insecticides (spinetoram and spinosad). If additional treatments are required after 2 consecutive applications of Group 5 insecticides, rotate to another class of effective insecticides for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area.

Note: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

TMTrademark of Dow AgroSciences LLC
R359-003
EPA accepted 08/13/09
Initial printing.