





FOR USE ON ASPARAGUS

EPA Reg. No. 241-418

Active Ingredient*:

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine	38.7%
Other Ingredients:	61.3%
Total:	100.0%

^{*1} gallon contains 3.8 pounds of pendimethalin formulated as an aqueous capsule suspension.

OBSERVE AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS, PRECAUTIONARY STATEMENTS, MIXING AND APPLICATION INSTRUCTIONS, AND FOLLOWCROP RESTRICTIONS ON THE Prowl® H_2O herbicide MAIN CONTAINER LABEL BEFORE USING.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label and the current EPA-accepted label for **Prowl H₂0**, **EPA Reg. No. 241-418**, must be in the possession of the user at the time of pesticide application.

Prowl H20 is a selective herbicide for controlling most annual grasses and certain broadleaf weeds as they germinate. For a list of weeds controlled, refer to the **Prowl H20** container label (see **Table 1**). Uniformly apply **Prowl H20** with properly calibrated ground equipment in 10 or more gallons of water, or in 5 or more gallons per acre of water by air. Use sprayers with constant agitation and equipped with nozzles that provide accurate and uniform application.

Adequate rainfall or overhead irrigation after application prior to weed seedling germination will provide the most effective weed control.

Always perform a mixing test to check the compatibility of **Prowl H₂O** with all potential tank mix partners.

Follow all precautions and restrictions on the labels of all products applied in combination with **Prowl H₂O**. Always follow the most restrictive label.

Endangered Species Protection

If endangered species occur in proximity to the application site, the following mitigation measures are required:

- If applied by ground, leave untreated buffer zone of 200 feet. The product must be applied using a low boom (20 inches above the ground) and ASAE fine to medium/coarse nozzles.
- If applied by air, leave an untreated buffer zone of 170 feet. Must use straight-stream nozzles (D-6 or larger); wind can be no more than 8 mph, and release height must be 15 feet or less.

To determine whether your county has an endangered species, consult the website http://www.epa.gov/espp/usa-map.htm.

Endangered Species Bulletins may also be obtained from extension offices or state pesticide agencies. If the bulletin is not available for your specific area, check with the appropriate local state agency to determine if known populations of endangered species occur in the area to be treated.

Spray Drift

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment-related and weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. It is the responsibility of the applicator to avoid spray drift onto nontarget areas.

Application Instructions

Prowl® H₂O herbicide may be applied by ground or air.

With a single application, uniformly apply **Prowl H₂O** to asparagus up to 8.2 pints per acre as a broadcast spray to the soil surface at least 14 days before the first harvest. Application must be made prior to spear emergence or remove emerged spears prior to making the application. If asparagus is grown on sandy soils, **DO NOT** apply **Prowl H₂O** at more than 2.4 pints per acre.

Prowl H_2O should be applied prior to weed germination. **Emerged weeds will not be controlled by this treatment.**

Prowl H₂O may be used on peat and muck soils, but weed control may be inconsistent and/or reduced. Use maximum labeled use rate allowed in asparagus.

Restrictions and Limitations

- **DO NOT** apply postemergence over the top of emerged spears as severe injury may occur.
- DO NOT apply more than 8.2 pints per acre per season.
- DO NOT apply within 14 days of harvest.
- DO NOT feed forage or graze livestock in treated fields.
- DO NOT apply by chemigation methods.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Prowl is a registered trademark of BASF.

© 2009 BASF Corporation. All rights reserved.

000241-00418.20090813**.NVA 2009-04-195-0153**

Based on: NVA 2008-04-195-0213

BASF Corporation 26 Davis Drive Research Triangle Park, NC 27709



The Chemical Company