

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Roundup PRO® Concentrate herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



EPA Reg. No. 524-529

Roundup PRO is a trademark of Monsanto Technology LLC.

For Hollow Stem Injection

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup PRO Concentrate herbicide for essential product performance information.

This product may be applied through hand-held injection devices that deliver recommended amounts of this product into targeted hollow-stem plants growing in any non-crop site specified on this label. For control of the following hollow-stem plants, follow the use instructions below:

Japanese Knotweed, *Polygonum cuspidatum*

Inject 5 mL/stem of this product between second and third internode.

Bohemian Knotweed, *Polygonum bohemicum*

Inject 5 mL/stem of this product between second and third internode.

Giant Hogweed, *Heracleum mantegazzianum*

Inject one leaf cane per plant 12 inches above root crown with 5 mL of a 5% v/v solution of this product.

Poison Hemlock, *Conium maculatum*

Inject one leaf cane per plant 10 to 12 inches above root crown with 5 mL of a 5% v/v solution of this product.

Field horsetail, *Equisetum arvense*

Inject one segment above the root crown with 0.5 mL/stem of this product. Use a small syringe that calibrates to this rate.

Canada Thistle, *Cirsium arvense*

Cut 8 to 9 of the tallest plants at bud stage in a clump with clippers. Use a cavity needle that is pushed into the stem center and then slowly removed as 0.5 mL/stem of this product is injected into the stem.

NOTE: Based on the maximum annual use rate of glyphosate for these non-crop sites, the combined total for all treatments must not exceed 8.5 quarts of this product per acre. At 5 mL per stem, 8.5 quarts should treat approximately 1600 stems.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup PRO Concentrate herbicide before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

© 2006 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

5/18/2006

21203E1-22